

Continuing Education: Improving Athletic Trainers' Knowledge of Musculoskeletal Diagnostic Ultrasound Through a Physician-Led Longitudinal Ultrasound Course

Allison N. Schroeder, MD*; Wade O. Johnson, MD†; Elena J. Jelsing, MD*‡; Amanda L. Williams, LAT, ATC‡; Joshua J. Pinkney, LAT, ATC‡; Chad A. Asplund, MD, MPH§; Ryan C. Kruse, MD||

*Department of Physical Medicine and Rehabilitation, Mayo Clinic, Rochester, MN; †Department of Physical Medicine and Rehabilitation, Mayo Clinic Health System, Mankato, MN; ‡Department of Orthopedic Surgery, Mayo Clinic, Minneapolis, MN; §Department of Family Medicine and Orthopedics, Mayo Clinic, Rochester, MN; ||Department of Orthopedics and Rehabilitation, University of Iowa Sports Medicine, Iowa City

Context: Musculoskeletal (MSK) diagnostic ultrasound (US) is increasingly being utilized by physicians to evaluate athletes in outpatient clinics, on the sidelines, and in athletic training facilities. Having a thorough understanding of MSK US will help athletic trainers (ATs) better assist physicians who perform MSK US.

Objective: To determine if a longitudinal 6-month MSK anatomy and US course would increase ATs' ability to acquire, label, and optimize US images of normal structures and improve their overall confidence in assisting with performing US and to determine if demographic factors, work-related factors, or higher self-reported confidence was associated with better performance.

Design: Prospective cohort study.

Setting: Academic institution.

Other Participants: Seventeen ATs working with sports medicine physicians at 3 institutions.

Interventions: Longitudinal 6-month MSK anatomy and US course.

Main Outcome Measures: Ultrasound image scores (normal structures scored on a 5-point scale) were assessed precourse, during the course, and postcourse. Scores were compared across time points for the assessed body regions (knee, ankle/foot, shoulder, elbow/wrist/hand, and hip). Associations with demographic factors, US experience, and AT self-reported confidence were explored.

Results: Seventeen ATs completed the course. There was a significant difference among the average overall precourse (average: 1.34/5, range: 0/5 to 4.23/5), during-course (average = 3.53/5; range, 2.6/5 to 4.37/5), and postcourse (average = 3.83/5; range, 2.33/5 to 4.67/5) image scores [$\chi^2(2) = 24.47$, $P < .001$]. There was a significant positive correlation between the numbers of days ($r_s[17] = 0.62$, $P = .01$) and hours ($r_s[17] = 0.55$, $P = .02$) per week that the AT spent observing or performing US scanning and the postcourse overall image scores. The ATs' confidence in identifying structures when they scan and postcourse overall image scores were marginally correlated but not significant ($r_s[17] = 0.47$, $P = .06$).

Conclusions: A comprehensive longitudinal MSK diagnostic US course may have resulted in significant improvements in an AT's ability to acquire, label, and optimize US images of normal MSK structures that are commonly injured by active persons.

Key Words: athletic trainer, musculoskeletal ultrasound, sports ultrasound, ultrasound education

Dr Schroeder is currently an Assistant Professor in the Department of Orthopedic Surgery, Division of Sports Medicine at University Hospital and Case Western Reserve University in Cleveland, OH. Address correspondence to Ryan C. Kruse, MD, Department of Orthopedics and Rehabilitation, University of Iowa Sports Medicine, 2701 Prairie Meadow Drive, Iowa City 52246. ryan-kruse@uiowa.edu.

Full Citation:

Schroeder AN, Johnson WO, Jelsing EJ, Williams AL, Pinkney JJ, Asplund CA, Kruse RC. Continuing education: improving athletic trainers' knowledge of musculoskeletal diagnostic ultrasound through a physician-led longitudinal ultrasound course. *Athl Train Educ J*. 2023;18(4):234–242.

Continuing Education: Improving Athletic Trainers' Knowledge of Musculoskeletal Diagnostic Ultrasound Through a Physician-Led Longitudinal Ultrasound Course

Allison N. Schroeder, MD; Wade O. Johnson, MD; Elena J. Jelsing, MD; Amanda L. Williams, LAT, ATC; Joshua J. Pinkney, LAT, ATC; Chad A. Asplund, MD, MPH; Ryan C. Kruse, MD

KEY POINTS

- A 6-month longitudinal MSK diagnostic US course improved ATs' abilities to acquire, label, and optimize US images of normal MSK structures. This course can be recreated in a setting with a physician US expert and ATs who are interested in assisting the physician with US evaluations.
- Athletic trainers who complete a diagnostic US course should spend time working directly with a physician who performs US as part of their job since spending more days or hours per week doing this was associated with higher image scores in the postcourse assessment.
- Understanding the basics of MSK diagnostic US will allow ATs to assist physicians when they are performing diagnostic and interventional US in the clinic, in the athletic training facility, and on the sidelines of sporting events. Athletic trainers may also play a role in the emerging field of "tele-US."

INTRODUCTION

Ultrasound (US) is an imaging modality that is increasingly being utilized by physicians practicing orthopedics and sports medicine. Diagnostic US can be performed as part of a patient evaluation in an outpatient clinic, in an athletic training facility, or on the sidelines of sporting events. Ultrasound can be utilized for diagnostic purposes to evaluate the muscle, tendon, ligament, bone, and bursae or guide interventional procedures. Ultrasound is an operator-dependent skill that requires a comprehensive understanding of anatomy, training in image acquisition and optimization, and hands-on practice. Hands-on,¹⁻⁴ instructor-led⁵⁻⁷ US training has been shown to be superior to didactic or self-guided training alone. Ultrasound training has been shown to improve the accuracy of physical examination palpation skills.⁸⁻¹⁰ Athletic trainers (ATs) work closely with physicians, taking histories, performing physical examinations, and assisting with the diagnosis and treatment of acute and chronic musculoskeletal (MSK) conditions. At our institutions, ATs work in the clinic, in the athletic training facility, and on the sidelines, assisting physicians with the capture, labeling, and optimization of US images during diagnostic US examinations and interventional US-guided procedures. If a physician is unable to be physically present, such as during an away game, ATs may perform "tele-US," an emerging practice where US scanning is performed by a novice under the direction of an expert through video calls in real time.¹¹⁻¹⁵

Given that many ATs work with sports medicine physicians who utilize US in their practice, an instructional MSK diagnostic US course was developed, offering free Board of Certification continuing education credits for the ATs working at our institutions. Previous studies have not evaluated MSK US training or ATs' ability to acquire and optimize diagnostic images after taking part in an anatomy and US training program. The purposes of this study were to (1) determine if a longitudinal 6-month MSK

anatomy and US course would increase ATs' US image scores overall and by body region (to assess their ability to acquire, label, and optimize US images of normal structures that may commonly be injured by active persons) and increase their overall confidence in performing US, (2) determine if higher US image scores were correlated with higher self-reported confidence, and (3) determine if demographic factors such as AT practice location and practice setting or work-related factors such as years practicing as an AT, years performing US, days per week performing or observing US, hours per week observing or performing US, or previous formalized US training were associated with higher US image scores. We hypothesized that the course would result in improvements in ATs' confidence and US image scores, that higher confidence would be correlated with higher image scores, and that weekly time spent observing or performing US would be associated with higher image scores.

METHODS

Study Design and Participants

Physicians invited all ATs with whom they worked (18 ATs) to participate in an MSK diagnostic US course for ATs that was conducted at the Mayo Clinic, Mayo Clinic Health Systems, and the University of Iowa. Seventeen ATs (10.8 ± 8.1 years of experience working as a certified AT) participated in the course. The ATs' demographics are summarized in Table 1. We collected precourse, during-course, and postcourse self-reported confidence scores and US images for those who consented to participate in the prospective cohort study, which was approved by the Mayo Clinic (no. 20-008617-02) and the University of Iowa Institutional Review Board (no. 202008451).

Musculoskeletal Diagnostic US Course

Athletic trainers participated in an MSK diagnostic US course that met weekly for 1-hour sessions (virtual didactics with hands-on sessions that were primarily in person, with an occasional virtual session^{16,17} due to participant mandatory quarantine secondary to the severe acute respiratory syndrome coronavirus 2 [SARS-CoV-2] pandemic, which occurred for 2 ATs for 1 week throughout the course) from November 2, 2020, through April 9, 2021. The course was taught by physicians trained in physical medicine and rehabilitation and sports medicine; the student-to-teacher ratio for the hands-on sessions was 2:1 to 4:1, depending on the site. A detailed timeline for the course is outlined in Figure 1.

Participant Self-Assessment Surveys

Participants were asked to complete precourse surveys administered in REDCap (Research Electronic Data Capture)^{18,19} assessing their years of experience as an AT; previous experience observing or performing US; weekly time spent observing or performing US in their current job; previous formal MSK US

Table 1. Baseline Athletic Trainer

Parameter	No. of ATs
Practice location	
University of Iowa	5
Mayo Clinic Health Systems	5
Mayo Clinic Square	7
Practice setting	
Clinical	10
Outreach	6
Other	1
Years observing MSK US	
<1	8
1.1–2	6
2.1–3	0
3.1–4	4
4.1–5	0
>5	6
Years of hands-on MSK US	
<1	14
1.1–2	1
2.1–3	2
>3	0
Days/week observing MSK US scanning	
0	5
1	5
2	1
3	3
4	1
5	1
Hours/week observing MSK US scanning	
<5	12
5–9.9	1
10–14.9	1
15–19.9	0
>20	3
Previous formalized US training	
No previous formal training	8
Lectures on MSK US but no hands-on practice	4
Guided hands-on practice	5

Abbreviations: ATs, athletic trainers; MSK, musculoskeletal; US, ultrasound.

instruction; confidence on a 0 to 100 scale in identifying structures when a physician is scanning, identifying structures when an AT (AT completing the survey) is scanning, labeling the US image, and optimizing the US image (depth/gain); and demographic information (see Supplemental File 1, available online at www.nataej.org). After each weekly session during the course, participants were asked to complete a survey confirming attendance for anatomy, US video viewing, and hands-on scanning (see Supplemental File 2). The same questions assessing AT confidence were repeated weekly during the course to assess the ATs' confidence for each body region and after the course to assess their overall confidence (see Supplemental File 3).

Image Acquisition, Review, and Scoring

Athletic trainers acquired specific images designated on the image structure list (Table 2) that was developed by the physicians and included 6 structures for each body region that are commonly encountered in sports medicine settings. Athletic

Figure 1. Athletic trainer MSK US course timeline. The MSK diagnostic US course began with an Introduction to US lecture and participants completing their precourse image acquisition. Thereafter, over a 4-week period, ATs attended an anatomy lecture (virtual, week 1), watched AMSSM (virtual) US videos (week 2), attended a live, hands-on, physician-led (E.J.J., W.O.J., R.C.K., and A.N.S.) scanning session (week 3), and obtained images for the studied body region (week 4). This was repeated for each of the following body regions: knee, ankle/foot, shoulder, elbow/wrist/hand, and hip. At the end of the course, ATs were given 6 months to collect postcourse images. Abbreviations: AMSSM, American Medical Society for Sports Medicine; MSK, musculoskeletal; US, ultrasound.



Table 2. Image Structure List Showing the Details of Anatomic Structures Imaged by the Athletic Trainers

Body Region	Structure of Interest—Surrounding Structure(s) (to Label)
Knee	<ul style="list-style-type: none"> • Patellar tendon just distal to the patella (short axis)—Hoffa's fat pad • Patellar tendon at the tibial tuberosity attachment (long axis)—deep infrapatellar bursa, Hoffa's fat pad • Suprapatellar recess (long axis)—quadriceps tendon in the long axis, prefemoral fat pad • Rectus femoris muscle at the mid thigh (short axis)—vastus medialis, vastus lateralis, vastus intermedius • Quadriceps tendon attaching to the patella (long axis)—patella, quadriceps fat pad • MCL (long axis)—medial meniscus, femur, tibia
Ankle/foot	<ul style="list-style-type: none"> • ATFL (long axis)—tibia, talus • AITFL (long axis)—tibia, fibula • Peroneus brevis insertion on base of the 5th metatarsal (long axis)—base of the 5th metatarsal • Central cord of the plantar fascia (long axis)—calcaneus • Achilles tendon (long axis)—calcaneus, Kager fat pad, retrocalcaneal bursa • Tibialis posterior tendon in the retromalleolar groove (short axis)—medial malleolus, tibial nerve
Shoulder	<ul style="list-style-type: none"> • Long head of the biceps tendon in the bicipital groove (long axis)—humerus • Long head of the biceps tendon in the bicipital groove (short axis)—greater tuberosity, lesser tuberosity • Supraspinatus tendon (long axis)—greater tuberosity, subacromial/subdeltoid bursa • Supraspinatus tendon (short axis)—greater tuberosity, deltoid • AC joint (short axis with the acromion and clavicle in the long axis)—acromion, clavicle • Posterior glenohumeral joint (with the labrum in view)—humeral head, glenoid, labrum
Elbow/wrist/ hand	<ul style="list-style-type: none"> • UCL (long axis)—ulna, medial epicondyle • Common extensor tendon (long axis)—lateral epicondyle • Ulnar nerve (short axis)—olecranon, medial epicondyle • Median nerve in the carpal tunnel (short axis, in the carpal tunnel inlet)—scaphoid, pisiform, transverse carpal ligament • Lister's tubercle—extensor pollicis longus in the 3rd dorsal compartment, extensor carpi radialis longus and brevis in the 2nd dorsal compartment (short axis) • 1st CMC joint (short axis with the bones in the long axis)—trapezium, 1st metacarpal
Hip	<ul style="list-style-type: none"> • Hip joint (view of the femoral head/neck in the long axis to the femoral shaft)—femoral head, femoral neck, joint capsule • AIIS (short axis)—rectus femoris tendon (in the short axis) • Subgluteus maximus/trochanteric (gluteus minimus in the oblique short axis)—greater trochanter anterior facet, greater trochanter lateral facet • Gluteus medius inserting on the greater trochanter (long axis)—lateral facet of the greater trochanter • Ischial tuberosity (short axis)—conjoint tendon, semimembranosus tendon • Iliopsoas tendon (short axis)—acetabulum, psoas muscle

Abbreviations: AC, acromioclavicular; AIIS, anterior inferior iliac spine; AITFL, anterior inferior tibiofibular ligament; ATFL, anterior talofibular ligament; CMC, carpometacarpal; MCL, medial collateral ligament; UCL, ulnar collateral ligament of the elbow.











trainers obtained all images of the structure precourse and postcourse. Each week during the course, ATs obtained images of structures for that specific body region (Figure 1). Participants were made aware of the physician-selected key components of US image acquisition and interpretation, included in the image scoring criteria, and were presented with examples of optimal and suboptimal image acquisition (Figure 2). Three sports medicine physicians who completed a physical medicine and rehabilitation residency (at different institutions) and a sports medicine fellowship (2 at the same institution with a different fellowship program director and 1 at a different institution) with a focus on diagnostic and interventional MSK US (W.O.J., R.C.K., and A.N.S.) independently evaluated deidentified images of normal structures obtained by the ATs. Since there is no previously validated system to score MSK US images, a scoring system was created by the study authors. Images were scored, with 0 = absent and 1 = present, for the following criteria: structure label, centered images, surrounding structure label, directionality indicated (eg, medial or lateral), and quality of the overall image (correction of anisotropy and optimization of depth and gain). The consensus value (ie, needed 2 reviewers to agree) for each criterion was used in the analysis. Each image was given a score out of a maximum of 5 points. These scores were then averaged

among all participants for each structure. An average score was calculated precourse, during the course, and postcourse for each body region and overall.

Statistical Analysis

Demographic data were reported using standard descriptive statistics. We calculated Krippendorff α to assess the interrater reliability between US image scorers (W.O.J., R.C.K., and A.N.S.). Before the study, one set of sample images was scored with consistency among all scorers. Pre-, during-, and postcourse confidence values were compared using a repeated-measures analysis of variance (ANOVA) with an unstructured covariance matrix; if time was significant, post hoc tests with a Tukey correction were used to identify significant differences in least-square means at each time point. Athletic trainer image scores were compared pre-, during-, and postcourse overall and by body regions using Friedman's ANOVA with a Tukey adjustment for post hoc comparisons. Spearman correlations were used to assess the correlation between AT confidence in identifying an image when the AT was scanning and their overall image scores pre-, during, and postcourse. Spearman correlations were also used to assess the correlation between AT overall postcourse image scores


Figure 2. Ultrasound image scoring criteria.

Scoring Criteria	Present/Optimized (scored 1)	Absent/Not Optimized (scored 0)
Structure Label: <ul style="list-style-type: none">The structure of interest is accurately labeled (with an arrow pointing to the structure, label right over the structure, or label at the bottom of the image)		
Centered: <ul style="list-style-type: none">The structure of interest is in the center of the screen and optimizedMiddle 1/3		
Surrounding structures: <ul style="list-style-type: none">The surrounding structures are accurately labeled (using arrows to point to each structure)		
Directionality: <ul style="list-style-type: none">The directionality of the transducer is accurately labeled (i.e. medial/lateral/proximal/distal)		
Optimization: <ul style="list-style-type: none">The depth is properly set for the image optimization (middle 1/3)Eliminate anisotropyImage is not oblique to structure of interest		

and the number of years working as an AT, the number of years spent observing US, the number of years learning or performing US scanning, and the number of days and hours per week that the AT spent observing or performing US scanning. Due to the

categorical nature of the variables, Kruskal-Wallis tests were used to determine the relationship of ATs' overall postcourse image scores with the study site, AT role, and formal US training. If a significant association was identified, pairwise comparisons were

Figure 3. Average AT confidence pre-, during, and postcourse. Athletic trainer confidence (out of 100) in identifying structures when the physician is scanning, identifying structures when the AT is scanning, their ability to properly label a structure, and their ability to adjust depth and gain is displayed. Chi-square and *P* values are also displayed. Abbreviation: AT, athletic trainer. *** indicates statistical significance (*P* < .05). Error bars represent standard deviations.



assessed, and *P* values were adjusted for type I errors using the Dwass-Steel-Critchlow-Flinger method.

RESULTS

Course Completion

All ATs reviewed the comprehensive anatomy lecture and the American Medical Society for Sports Medicine (AMSSM) lectures for each body region, except for 1 participant who did not review the comprehensive anatomy lecture for elbow/wrist/hand or watch the AMSSM videos for hip. All participants attended the hands-on sessions for each body region. For each of the 17 participants, a total of 90 images were scored by each of the 3 image scorers (W.O.J., R.C.K., and A.N.S.). The interrater reliability of image scoring was relatively high (Krippendorff α = 0.71, 95% confidence interval [CI] = 0.69, 0.73).

Athletic Trainer Confidence

A significant difference in AT confidence pre-, during, and postcourse (Figure 3) was detected for identifying a structure when the physician was scanning, properly labeling an image, and adjusting depth/gain (all *P* < .001). No significant difference was detected in identifying a structure when the AT was scanning (*P* = .30). Post hoc comparisons revealed significant pairwise differences between all time points for confidence in identifying a structure when the physician was scanning (*P* values pre- to during course of < .001, pre- to postcourse of < .001, and during to postcourse of .03). Confidence in labeling an image and confidence in optimizing depth/gain pre- to during course, pre- to postcourse, and during to precourse were statistically significant (during to postcourse *P* = .04 for labeling; all other *P* < .001).

Athletic Trainer US Image Scores

Overall, we found that the ATs' image scores improved throughout the course from 1.34/5 precourse (range, 0/5 to 4.23/5), to

3.53/5 during the course (range, 2.6/5 to 4.37/5), to 3.83/5 postcourse (range, 2.33/5 to 4.67/5). There was a significant difference among the average precourse, during-course, and postcourse image scores overall ($\chi^2[2] = 24.47$, *P* < .001) and for each body region (all *P* < .01) (Table 3). Post hoc testing revealed that for knee and shoulder, all time points were significantly different (all *P* < .01). For ankle/foot, all time points were significantly different (*P* < .01 for all except during versus postcourse [*P* = .04]). For elbow and hip, pre- versus during course and pre- versus postcourse were significantly different (all *P* < .01), but during versus postcourse was not significantly different (*P* = .22 for elbow and *P* = .99 for hip). The highest scores were recorded for the knee (precourse = 1.94/5, during course = 3.98/5, and postcourse = 4.44/5), and the lowest scores were recorded for the hip (precourse = 0.85/5, during course = 3.23/5, and postcourse = 3.20/5) at all time points. The average postcourse image scores for each anatomic structure are displayed in Figure 4.

Association of AT Characteristics and Confidence With US Image Scores

We found a positive correlation between ATs' confidence in their ability to identify US structures when the AT was scanning and overall image scores. This was statistically significant precourse ($r_s[17] = 0.64$, *P* = .005) but was only marginally correlated but not significant postcourse ($r_s[17] = 0.47$, *P* = .06). During the course, there was a significant positive correlation between ATs' confidence in their ability to identify US structures when the AT was scanning and image scores for the ankle/foot ($r_s[17] = 0.57$, *P* = .02) and shoulder ($r_s[17] = 0.55$, *P* = .02). No significant correlations between ATs' confidence in their ability to identify US structures when the AT was scanning and image scores for the knee ($r_s[17] = -0.17$, *P* = .95), elbow/wrist/hand ($r_s[17] = 0.44$, *P* = .08), and hip ($r_s[17] = -0.15$, *P* = .57) were identified.

No significant associations between the average score of all images collected during postcourse and study site (*P* = .47), years working as an AT (*P* = .62), AT role (clinical, outreach,

Table 3. Athletic Trainer Ultrasound Average Image Scores for Each Body Region

Body region	Precourse Average \pm SE	During-Course Average \pm SE	Postcourse Average \pm SE	Freidman Q (χ^2) Among Time Points	P Value Among Time Points
Knee	1.94 \pm 1.46	3.98 \pm 0.58	4.44 \pm 0.44	28.22	<.001
Ankle/foot	1.26 \pm 1.37	3.69 \pm 0.49	3.97 \pm 0.60	26.06	<.001
Shoulder	1.37 \pm 1.62	3.47 \pm 0.69	3.91 \pm 0.72	24.27	<.001
Elbow/wrist/hand	1.28 \pm 1.36	3.27 \pm 1.09	3.63 \pm 1.02	23.47	<.001
Hip	0.85 \pm 1.25	3.23 \pm 0.86	3.20 \pm 0.96	28.35	<.001
Total	1.34 \pm 1.31	3.53 \pm 0.51	3.83 \pm 0.61	24.27	<.001

Abbreviation: SE, standard error.

or other) ($P = .44$), the number of years spent observing US before the course ($P = .49$), the number of years learning or performing hands-on scanning ($P = .52$), or US instruction before this course (none, didactic only, or hands-on scanning) ($P = .79$) were detected. There was a significant positive association of the number of days per week that the AT currently spent observing or performing US scanning ($P = .008$) and the number of hours per week that the AT currently spent observing or performing US scanning ($P = .02$) with the postcourse overall image scores, although both groups of ATs who spent time observing or performing US weekly ($n = 6$, average overall scores of 2.36 precourse and 4.25 postcourse) and ATs who did not spend time observing or performing US weekly ($n = 11$, average overall scores of 0.43 precourse and 3.61 postcourse) improved throughout the course.


DISCUSSION

Our results showed that during this 6-month, longitudinal MSK diagnostic US course, AT US image scores significantly improved over time. Although slight, the postcourse image scores were higher than the during-course image scores, indicating that the improvements made between precourse image acquisition and image acquisition during the course were retained, and even

slightly increased, over time. This is likely due to the regular repeated assessments and longitudinal approach to the course as retention of acquired knowledge and skills are enhanced with continued exposure to content and iterative practice with more time with hands on the transducer.^{20–22} Additionally, ATs who spent more days and more hours per week observing or performing US with a physician had higher scores on the postcourse images. This underscores the importance of iterative practice and exposure to US with frequent exposure to a physician US expert for on-the-job learning. Overall, the MSK diagnostic US course was successful in enhancing the ATs' abilities to acquire, label, and optimize US images of normal structures, but having continual exposure to US as part of their job is significantly important in enhancing the ATs' US abilities.

The ATs' confidence in their ability to identify structures when an AT was scanning did not increase throughout the course, but their confidence in their ability to identify structures when the physician was scanning, their ability to properly label, and their ability to optimize depth and gain did increase. This may be due to the fact that most ATs working with a physician identify images while the physician is scanning, label images, and optimize depth and gain on a regular basis, but they rarely perform US scanning in their daily practice. Previous studies of physicians have revealed a correlation between confidence and US

Figure 4. Average postcourse scores for each structure. Structures are noted on the x-axis, and the image score out of 5 points is noted on the y-axis. Abbreviations: AC, acromioclavicular; AIIS, anterior inferior iliac spine; AITFL, anterior inferior tibiofibular ligament; AT, athletic trainer; ATFL, anterior talofibular ligament; CMC, carpometacarpal; LAX, long axis; MCL, medial collateral ligament; SAX, short axis; UCL, ulnar collateral ligament of the elbow.



skill level, but one study assessed the interpretation, not the acquisition, of images by emergency medicine residents and physicians²³; one reported improvements in identifying normal images of the wrist but did not evaluate image optimization by radiology residents and MSK fellows²⁴; and another assessed performing and interpreting abnormal focused abdominal US examinations.²⁵ In our study, ATs were acquiring images of normal structures and were asked to optimize images rather than identify pathology. Overall, this MSK diagnostic US course improved ATs' confidence in performing the tasks that they more commonly perform in our clinics and athletic training facilities: identifying structures when the physician is scanning, labeling images, and optimizing depth and gain. However, self-reported confidence in their ability to acquire or identify US images when they are scanning should be interpreted cautiously, as this did not correlate with the US image acquisition skill level.

Strengths and Limitations

There were several strengths to this study. This study evaluated a comprehensive, 6-month longitudinal MSK diagnostic US course that covered peripheral joints and surrounding soft tissue structures. All participants attended the hands-on scanning sessions, completed surveys, and submitted US images throughout the course. Previously, the effects of US training on a learner's ability to acquire and optimize diagnostic US images have been limited in scope to isolated body regions^{26,27} or typically evaluated for resident or fellow physicians.²⁴ To our knowledge, this was the first study to assess a learner's ability to acquire US images of normal MSK structures and can serve as a model for future studies that have MSK US image acquisition as an outcome measure.

There were also several limitations to this study. This study encompassed only a single year, and it is unclear if the information learned during the course will be retained over subsequent years. Implementing a similar course at other institutions may present a challenge, as a physician skilled in MSK US is necessary to lead the course. It is unclear if a different US curriculum would produce similar results. In addition, completion of the course required a weekly commitment by the ATs at our institutions, which they did outside their typical work hours. Athletic trainers spent roughly 1 hour per week reviewing the anatomy lectures and the AMSSM US videos and roughly 2 to 3 hours per week during the hands-on session and acquiring US images. Obtaining pre- and postcourse images took several hours and, due to busy sports seasons, took some ATs the full 6 months to complete the image acquisition as part of the postcourse assessment. Finally, the overall improvement in posttest image scoring may have been influenced by factors outside the course, such as the time of exposure to US in clinical settings; however, given that ATs who observed and performed US while working with a physician on a weekly basis and those who did not both improved throughout the course, participation in the course likely contributed to the overall improvement in image acquisition scores that was observed.

The subjective image scoring system that we employed had not been previously described or validated; however, this system was developed to assess the key components of MSK US image acquisition and optimization. Measures to decrease the subjective nature of the scoring system were employed, including obtaining consensus among 2/3 of the image reviewers and having well-defined criteria. However, there was relatively high

interrater reliability between the physician scorers for each, and future studies assessing learners' ability to acquire MSK US images could utilize this scoring system. Additionally, the ability to acquire US images of normal structures was assessed, and any correlation with the ability to acquire US images of pathologic structures is unknown. Finally, although the image scoring system was utilized to assess key components of US image acquisition and optimization, it was not designed to determine criteria that would indicate that an AT is competent in performing US under a physician's guidance.

CONCLUSIONS

Ultrasound is an imaging modality that is increasingly being utilized by physicians and the ATs who work with them in sports medicine clinics, in athletic training facilities, and on the sidelines. We have described a comprehensive longitudinal MSK diagnostic US course and found that participation may have resulted in significant improvements in ATs' abilities to acquire, label, and optimize US images of normal MSK structures that are commonly injured by active persons. Athletic trainers who spend more time with physicians who utilize US had the best image scores. Developing a thorough understanding of MSK US will allow ATs to better assist physicians when they are performing diagnostic and interventional US in the clinic, in the athletic training facility, and on the sidelines of sporting events.

ACKNOWLEDGMENTS

We acknowledge Jeffrey M. Payne, MD, and Shelby Johnson, MD, for their assistance with teaching the hands-on portion of the ultrasound course and Stephanie Kliethermes, PhD, for her assistance in performing the statistical analysis.

REFERENCES

1. Wu WT, Chang KV, Han DS, Özçakar L. Musculoskeletal ultrasound workshops in postgraduate physician training: a pre- and post-workshop survey of 156 participants. *BMC Med Educ.* 2019;19(1):362.
2. Maw A, Jalali C, Jannat-Khah D, et al. Faculty development in point of care ultrasound for internists. *Med Educ Online.* 2016; 21:33287.
3. Greenstein YY, Littauer R, Narasimhan M, Mayo PH, Koenig SJ. Effectiveness of a critical care ultrasonography course. *Chest.* 2017;151(1):34–40.
4. Sarkar PK, Boivin M, Mayo PH. Effectiveness of an advanced critical care echocardiography course. *J Intensive Care Med.* 2020; 35(11):1332–1337.
5. Charnoff J, Naqvi U, Weaver M, Price C. Resident education of ultrasound-guided procedures: a homemade practice model pilot study. *Am J Phys Med Rehabil.* 2019;98(10):e116–e118.
6. Farjad Sultan S, Shorten G, Iohom G. Simulators for training in ultrasound guided procedures. *Med Ultrason.* 2013;15(2):125–131.
7. Micks T, Sue K, Rogers P. Barriers to point-of-care ultrasound use in rural emergency departments. *CJEM.* 2016;18(6):475–479.
8. Walrod BJ, Schroeder A, Conroy MJ, et al. Does ultrasound-enhanced instruction of musculoskeletal anatomy improve physical examination skills of first-year medical students? *J Ultrasound Med.* 2018;37(1):225–232.

9. Walrod BJ, Boucher LC, Conroy MJ, et al. Beyond bones: assessing whether ultrasound-aided instruction and practice improve unassisted soft tissue palpation skills of first-year medical students. *J Ultrasound Med*. 2019;38(8):2047–2055.
10. Rho ME, Chu SK, Yang A, Hameed F, Lin CY, Hurh PJ. Resident accuracy of joint line palpation using ultrasound verification. *PM R*. 2014;6(10):920–925.
11. Jensen SH, Weile J, Aagaard R, et al. Remote real-time supervision via tele-ultrasound in focused cardiac ultrasound: a single-blinded cluster randomized controlled trial. *Acta Anaesthesiol Scand*. 2019;63(3):403–409.
12. Salerno A, Tupchong K, Verceles AC, McCurdy MT. Point-of-care teleultrasound: a systematic review. *Telemed J E Health*. 2020;26(11):1314–1321.
13. Baribeau Y, Sharkey A, Chaudhary O, et al. Handheld point-of-care ultrasound probes: the new generation of POCUS. *J Cardiothorac Vasc Anesth*. 2020;34(11):3139–3145.
14. Kwon D, Bouffard JA, van Holsbeeck M, et al. Battling fire and ice: remote guidance ultrasound to diagnose injury on the International Space Station and the ice rink. *Am J Surg*. 2007;193(3):417–420.
15. Kirkpatrick AW, Blaivas M, Sargsyan AE, et al. Enabling the mission through trans-Atlantic remote mentored musculoskeletal ultrasound: case report of a portable hand-carried tele-ultrasound system for medical relief missions. *Telemed J E Health*. 2013;19(7):530–534.
16. Schroeder AN, Kruse RC. The future of virtual sports ultrasound education and collaboration. *Curr Sports Med Rep*. 2021;20(1):57–61.
17. Schroeder AN, Hall MM, Kruse RC. Sports ultrasound training during a pandemic: developing a “hands-on” skill through distance learning. *Am J Phys Med Rehabil*. 2020;99(9):860–862.
18. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42(2):377–381.
19. Harris PA, Taylor R, Minor BL, et al. The REDCap consortium: building an international community of software platform partners. *J Biomed Inform*. 2019;95:103208.
20. Xue G, Mei L, Chen C, Lu ZL, Poldrack R, Dong Q. Spaced learning enhances subsequent recognition memory by reducing neural repetition suppression. *J Cogn Neurosci*. 2011;23(7):1624–1633.
21. Larsen DP, Butler AC, Roediger HL III. Repeated testing improves long-term retention relative to repeated study: a randomised controlled trial. *Med Educ*. 2009;43(12):1174–1181.
22. Kerfoot BP, DeWolf WC, Masser BA, Church PA, Federman DD. Spaced education improves the retention of clinical knowledge by medical students: a randomised controlled trial. *Med Educ*. 2007;41(1):23–31.
23. Davis DP, Campbell CJ, Poste JC, Ma G. The association between operator confidence and accuracy of ultrasonography performed by novice emergency physicians. *J Emerg Med*. 2005;29(3):259–264.
24. Maetani TH, Schwartz C, Ward RJ, Nissman DB. Enhancement of musculoskeletal radiology resident education with the use of an individual smart portable ultrasound device (iSPUD). *Acad Radiol*. 2018;25(12):1659–1666.
25. Jang T, Naunheim R, Sineff S, Aubin C. Operator confidence correlates with more accurate abdominal ultrasounds by emergency medicine residents. *J Emerg Med*. 2007;33(2):175–179.
26. Swamy V, Brainin P, Biering-Sørensen T, Platz E. Ability of non-physicians to perform and interpret lung ultrasound: a systematic review. *Eur J Cardiovasc Nurs*. 2019;18(6):474–483.
27. Shah S, Santos N, Kisa R, et al. Efficacy of an ultrasound training program for nurse midwives to assess high-risk conditions at labor triage in rural Uganda. *PLoS One*. 2020;15(6):e0235269.