

Klykken LW, Pietrosimone BG, Kim KM, Ingersoll CD, Hertel J. Motor-neuron pool excitability of the lower leg muscles after acute lateral ankle sprain. *J Athl Train*. 2011;46(3):263–269.

Table 2 was printed incorrectly; please see the corrected table below. We regret the error.

Table 2. Maximal Hoffmann Reflex to Maximal Muscle Pooled Response Ratios Between Groups (Mean $\pm$ SD [95% Confidence Interval])

Muscle	Group			
	Acute Ankle Sprain		Injury Matched	
	Injured	Healthy	Uninjured	Healthy
Soleus	0.63 $\pm$ 0.23 (0.46, 0.80) ^a	0.47 $\pm$ 0.24 (0.08, 0.93)	0.58 $\pm$ 0.24 (0.42, 0.76)	0.54 $\pm$ 0.25 (0.37, 0.72)
Fibularis longus	0.22 $\pm$ 0.12 (0.14, 0.29)	0.25 $\pm$ 0.22 (0.12, 0.38)	0.24 $\pm$ 0.11 (0.17, 0.31)	0.24 $\pm$ 0.18 (0.13, 0.36)
Tibialis anterior	0.06 $\pm$ 0.07 (0.01, 0.10) ^b	0.22 $\pm$ 0.22 (0.09, 0.35)	0.15 $\pm$ 0.08 (0.1, 0.20)	0.18 $\pm$ 0.18 (0.07, 0.29)

^a Indicates greater than the uninjured limb ($P = .01$).

^b Indicates less than the uninjured limb ($P = .07$).