

# Reliability and Validity Evidence of Multiple Balance Assessments in Athletes With a Concussion

Nicholas Murray, PhD\*; Anthony Salvatore, PhD†; Douglas Powell, PhD‡; Rebecca Reed-Jones, PhD§

\*Department of Health and Kinesiology, Georgia Southern University, Statesboro; †The University of Texas at El Paso; ‡Campbell University, Buies Creek, NC; §University of Prince Edward Island, Charlottetown

**Context:** An estimated 300 000 sport-related concussion injuries occur in the United States annually. Approximately 30% of individuals with concussions experience balance disturbances. Common methods of balance assessment include the Clinical Test of Sensory Organization and Balance (CTSIB), the Sensory Organization Test (SOT), the Balance Error Scoring System (BESS), and the Romberg test; however, the National Collegiate Athletic Association recommended the Wii Fit as an alternative measure of balance in athletes with a concussion. A central concern regarding the implementation of the Wii Fit is whether it is reliable and valid for measuring balance disturbance in athletes with concussion.

**Objective:** To examine the reliability and validity evidence for the CTSIB, SOT, BESS, Romberg test, and Wii Fit for detecting balance disturbance in athletes with a concussion.

**Data Sources:** Literature considered for review included publications with reliability and validity data for the assessments of balance (CTSIB, SOT, BESS, Romberg test, and Wii Fit) from PubMed, PsycINFO, and CINAHL.

**Data Extraction:** We identified 63 relevant articles for consideration in the review. Of the 63 articles, 28 were considered appropriate for inclusion and 35 were excluded.

**Data Synthesis:** No current reliability or validity information supports the use of the CTSIB, SOT, Romberg test, or Wii Fit for balance assessment in athletes with a concussion. The BESS demonstrated moderate to high reliability (interclass correlation coefficient = 0.87) and low to moderate validity (sensitivity = 34%, specificity = 87%). However, the Romberg test and Wii Fit have been shown to be reliable tools in the assessment of balance in Parkinson patients.

**Conclusions:** The BESS can evaluate balance problems after a concussion. However, it lacks the ability to detect balance problems after the third day of recovery. Further investigation is needed to establish the use of the CTSIB, SOT, Romberg test, and Wii Fit for assessing balance in athletes with concussions.

**Key Words:** Clinical Test of Sensory Organization and Balance, Sensory Organization Test, Balance Error Scoring System, Romberg test, Wii Fit, sensitivity, specificity

## Key Points

- No current reliability or validity data exist to support the use of the Wii Fit to diagnose balance impairments in athletes with concussions.
- The Balance Error Scoring System has moderate to high reliability and low to moderate validity evidence to support its use in diagnosing balance impairment in athletes with concussions.

An estimated 300 000 sport-related concussion injuries occur in the United States per year.<sup>1</sup> The rate of sport-related concussion is anticipated to rise as sports involvement increases at the collegiate and high school levels.<sup>2</sup> Concussion has been defined by the Third International Conference on Concussion in Sport as a complex pathophysiologic process affecting the brain induced by traumatic biomechanical forces.<sup>3,4</sup> Potential concussion symptoms include confusion, loss of memory, loss of consciousness, loss of spatial and temporal awareness, headaches, migraines, speech impairment, dizziness, nausea, balance disturbances, oculomotor control reduction, vision impairment, speech reduction, gait unsteadiness, and poor coordination.<sup>3</sup> Because of the variable symptoms experienced by affected athletes and the multiple mechanisms of injury that can contribute to a concussive injury, no 2 concussions are perfectly similar.<sup>3</sup> This hinders accurate assessment and management of sport-related concussion.

No single tool can fully quantify a concussion injury.<sup>5</sup> Yet measures that focus on the most prevalent symptoms experienced by individuals after a concussion injury could help in diagnosis. The most common symptoms are headaches, dizziness, and balance problems.<sup>5,6</sup> Of the 3 top symptoms, dizziness and balance dysfunction are related. Thus, examining balance dysfunction after a concussion could provide a quantitative, objective measure of the injury.

Of individuals suffering from a sport-related concussion, 30% report balance dysfunction and 75.6% report dizziness as a debilitating symptom.<sup>7,8</sup> Balance disturbances have been noted to return to normal within 72 hours; however, prolonged damage may last more than 7 days beyond the initial injury.<sup>6,9</sup> Balance disturbance is the inability to stand with an upright posture without deviating outside the limits of the base of support.<sup>6</sup> After concussion, problems with the vestibular system are considered most likely to be responsible for the individual's inability to maintain balance.<sup>6</sup> Two candidate

mechanisms have been proposed to underlie diminished vestibular function with concussion: (1) damage to peripheral receptors or (2) inhibited sensory integration in response to structural damage of the central processing structures.<sup>6</sup> Vestibular dysfunction is a considerable detriment to activities of daily living and athletics. A balance disturbance could place the athlete at greater risk for additional injury through falls or collisions. Therefore, we need to accurately assess balance in athletes with concussions.

## CURRENT CLINICAL BALANCE ASSESSMENTS

Multiple tests have been developed to assess balance dysfunction in individuals with concussions. Balance assessments range from simple clinical sideline tests to complex laboratory testing. The most common assessments discussed in the concussion literature are the Clinical Test of Sensory Organization and Balance (CTSIB), the Sensory Organization Test (SOT), the Balance Error Scoring System (BESS), and the Romberg test or scale.<sup>6,10</sup> The CTSIB, BESS, and Romberg test are low-technology assessments of balance that are subjective and rely upon trained rater interpretations of balance deficits.<sup>6</sup> The SOT uses laboratory-grade equipment to assess balance performance. However, the SOT is not cost-effective or easily accessible to the average clinician.<sup>6,11</sup> Because of limitations in the current balance assessments, the National Collegiate Athletic Association has adopted the Wii Fit (Nintendo, Redmond, WA) as an alternative method to assess postural control after a possible concussion and during the return-to-play process.<sup>12</sup> Yet limited evidence exists to support the reliability and validity of the Wii Fit as a balance-assessment instrument in athletes with concussions. Also, limited reliability and validity evidence supports the use of the other balance-assessment instruments in individuals with concussions.

Therefore, the purpose of our systematic review was to examine the reliability and validity evidence for the Wii Fit balance game compared with traditional balance assessment by the CTSIB, the SOT, the BESS, and the Romberg test in athletes with concussions.

## METHODS

### Reliability and Validity Definitions

*Reliability* is the ability of a test to demonstrate the same results over time.<sup>13</sup> Reliability coefficients, specifically interrater and intrarater, that range from 0 to 1 (0 = *no reliability*, 1 = *perfect reliability*) are the most commonly reported methods.<sup>13</sup> For this review, we considered studies that assessed and contained reliability coefficients.

*Validity* refers to the ability of a test to be accurate and measure what it is supposed to measure.<sup>13</sup> There are many forms of validity, including diagnostic validity, which is critical in the evaluation of an injury or disease. Correct clinical diagnosis relies upon the ability of a test to indicate if the patient has or does not have the condition.<sup>13</sup> Sensitivity (true positive) and specificity (true negative) are the statistical indexes most commonly used for diagnostic validity of a test within the health sciences.<sup>13</sup> Therefore, we reviewed sensitivity and specificity measures for each test of balance dysfunction in individuals with concussions.

**Table 1. Inclusion and Exclusion Criteria for all Articles Reviewed**

Inclusion
History of balance assessment
Basic knowledge of balance assessment
Concussion research
Neurologic condition similar to concussion (Parkinson disease, traumatic brain injury)
Aging population
Reliability or validity evidence pertaining to assessment
Exclusion
Non-English articles
Sample size <10
Pilot study
Unpublished articles
Amputee research (lower, upper, or both)
Non-research-related articles
No reliability coefficient reported
No sensitivity or specificity coefficients reported
Qualitative study methods (surveys, etc)


### Current Clinical Balance Assessments

Articles relating to the history, reliability, and validity of the current clinical balance assessments were searched using the key words (1) *Clinical Test of Sensory Organization and Balance AND reliability OR validity OR sensitivity AND specificity*, (2) *CTSIB AND reliability OR validity OR sensitivity AND specificity*, (3) *CTSIB AND concussion*, (4) *Sensory Organization Test AND reliability OR validity OR sensitivity OR specificity*, (5) *SOT AND reliability OR validity OR sensitivity OR specificity*, (6) *SOT AND concussion*, (7) *Balance Error Scoring System AND reliability OR validity OR sensitivity OR specificity*, (8) *BESS AND reliability OR validity OR sensitivity OR specificity*, (9) *BESS AND concussion*, (10) *Romberg*, (11) *Romberg AND balance*, (12) *sharpened Romberg AND balance*, (13) *Romberg AND reliability*, (14) *Romberg AND validity*, and (15) *Romberg AND concussion*.

The literature search was limited to the dates January 1, 1990, to January 31, 2013. We included a publication if it contributed to the understanding of the CTSIB, SOT, BESS, or Romberg test and was used in concussion research or research on neurologic conditions similar to a concussion, specifically traumatic brain injury or Parkinson disease. A publication was excluded if it did not meet the inclusion criteria. Publications posted to PubMed, PsycINFO, and CINAHL that used the CTSIB, SOT, BESS, or Romberg test in balance studies and all reliability and validity evidence of each test were examined. We selected these databases because they reflect the major contributing literature in the allied health sciences without potential overlap of citations. Our search identified and retrieved 58 articles that satisfied the inclusion criteria and pertained to 1 of the balance assessments, with an emphasis on concussion, reliability, or validity (Table 1 and Figure 1).

### Wii Fit Balance Assessment

Articles related to the Wii Fit history, reliability, and validity were searched using the key words (1) *Wii Fit*, (2) *Wii Fit AND*, (3) *Wii Fit AND reliability*, (4) *Wii Fit AND validity*, and (5) *Wii Fit AND concussion*. The search was limited to the dates January 1, 1990, to January 31, 2013. We identified publications that addressed the use of the Wii




**Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2009 flow diagram of inclusion and exclusion of articles relating to the traditional balance measures.**

Fit in concussion research or research on a similar neurologic condition to a concussion. A publication was excluded if it did not meet the inclusion criteria. PubMed, PsycINFO, and CINAHL were reviewed for information regarding the use of the Wii Fit with balance or general reliability and validity data across all demographics. We included these databases because they comprise the major contributing literature of the allied health sciences without potential overlap. Our search revealed 14 articles that were related to the purposes of the study; 6 relevant articles from the search were retrieved and included in the review (Figure 2). Specific inclusion and exclusion criteria are found in Table 1.

## RESULTS

We identified 63 relevant articles for consideration in the review. Of the 63 articles, 28 were considered appropriate for inclusion in the review and 35 were excluded (Figures 1 and 2). Articles were rejected because they were duplicates, had no relationship to concussion or similar neurologic deficits (Methods section for exact criteria), or did not present statistically significant reliability or validity data. For example, some authors reported that the balance test was reliable (without coefficients) and valid (without sensitivity or specificity). We rejected these studies because of the lack of statistically significant reliability or validity data. Following the PRISMA 2009 guidelines<sup>14</sup> for including significant measurement results, study strength,



**Figure 2. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2009 flow diagram of inclusion and exclusion of articles relating to the Wii Fit.**

and weaknesses, we rated articles on an alphabetical scale (A–F). Articles rated A had a robust sample size (30+), had reliability and validity coefficients with appropriate magnitude calculations, provided power analysis, involved quantitative study methods, and were easy to read. Articles rated B had a sample size of  $\geq 15$ , supplied reliability and validity coefficients (lacking magnitude calculations), and were based on quantitative study methods. Articles rated C had a low sample size ( $< 15$ ), gave reliability and validity coefficients, or were questionnaire-based studies. Articles rated D lacked a sufficient sample size ( $\leq 15$ ), did not include reliability or validity coefficients, and involved qualitative study methods. Articles rated F were considered pilot studies ( $< 10$  sample size) or unpublished research and did not include reliability or validity coefficients.

Articles rated F were deemed inappropriate for inclusion and were excluded from this systematic review (Table 1). Articles with a rating of D were also rejected unless they contributed solely to the historical context of the tests in question; we included only 1 article with a D rating in this review. All of the articles included in this review are described in Table 2. The citations in the evidence table include the above-mentioned information in alphabetical order by the first author's last name. Opinion articles, consensus statements, handbooks, and other nonoriginal research included in the evidence table were not rated but were included in the study because they made a contribution to the development of the argument we present. All authors contributed to identifying and screening the retrieved articles and making recommendations as to which articles should be included; however, the first

**Table 2. Evidence Table for Included Articles as Rated by Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2009 Alphabetical Rating Scale Extended on Next Page**

Citation	Year	Purpose	No.	Synthesis
NCAA Handbook <sup>12</sup>	2012	NCAA support of methods of treatment for collegiate athletes	N/A	Supports use of Wii Fit in concussion management
Aubry et al <sup>13</sup>	2002	Summary of 1st international conference on concussion	N/A	Information on how to properly manage athletic concussion
Bell et al <sup>15</sup>	2011	Systematic review of BESS reliability and validity evidence in all clinical use	N/A	BESS reliability and validity vary greatly by populations
Cavanaugh et al <sup>16</sup>	2005	Support for return to play for individuals with concussions with use of COP, not Romberg	N/A	More precise to measure balance with COP over time theory rather than Romberg. Posture control is a multivariate process. Fewer or more organized DOF reduce adaptability. Changes in COP oscillations in PC reflect changes.
Clark et al <sup>17</sup>	2010	Static validity and reliability of the Wii Fit compared with force plates	30	ICC = 0.94 within device, 0.89 between Wii Fit and AMTI force plate
Ford-Smith et al <sup>18</sup>	1995	Test-retest reliability of the SOT in OA	40	Fair to good reliability for the SOT
Guskiewicz et al <sup>6</sup>	2011	Review of balance assessments for concussion	N/A	Balance disturbances only occur in approximately 30% (Romberg test; not appropriate for athletes)
Guskiewicz et al <sup>7</sup>	2007	Head impacts in collegiate football players and relationship to concussion	88	13 concussions reported in 88 participants
Guskiewicz et al <sup>19</sup>	1996	Examined the CTSIB over the recovery period of a concussion and its effects on postural stability	38	Athletes with concussions reported greater postural sway during balance assessment that was significantly different from matched controls
Holmes et al <sup>20</sup>	2013	Determine validity of Wii Fit balance board in static posture of individuals with Parkinson disease	20	Mean ICC of 0.96
Ingersoll and Armstrong <sup>21</sup>	1992	Effects of closed-head injury upon postural sway	48	First indication of Romberg being used to assess postural sway
Jacobson et al <sup>22</sup>	2011	Determine validity of modified Romberg test	103	Modified Romberg not sensitive enough to identify vestibular dysfunction
Khasnis and Gokula <sup>23</sup>	2003	Background info on Romberg	N/A	History of Romberg development and use
Lanska and Goetz <sup>24</sup>	2000	Background info on Romberg	N/A	History of Romberg development and use
McCorry et al <sup>4</sup>	2005	Consensus statement	N/A	Consensus statement on the areas of concussion from the Prague statement
McCrea et al <sup>25</sup>	2003	Acute effects and recovery time of collegiate concussions	1631	Examination of recovery time and return-to-play decisions in collegiate football
Michalski et al <sup>26</sup>	2012	Determine if Wii Fit games use different motor-control strategies	16	After 10 trials, specific learning effect is noted
Reed-Jones et al <sup>27</sup>	2012	How does Wii Fit balance game relate to current standardized tests of balance?	34	Wii Fit results do not correlate with standardized functional balance assessments
Riemann and Guskiewicz <sup>10</sup>	2000	Assessment of postural stability concerning mild head injury	32	BESS is useful clinical tool to assess balance dysfunction in athletes with concussions
Shumway-Cook and Horak <sup>28</sup>	1986	Assessment of sensory interaction of balance		Creation of CTSIB
Steffen and Seney <sup>29</sup>	2008	Examination of different balance measurements in individuals with Parkinson disease	37	Sharpened Romberg test is a good assessment of balance in individuals with Parkinson disease
Whitney et al <sup>30</sup>	1998	Review of balance assessments in clinical use in OA	N/A	CTSIB measures narrow aspects of balance
Wikstrom <sup>31</sup>	2012	Determine validity of balance scores produced by Wii Fit and reliability intrasession/ intersession	45	Poor reliability between Wii Fit and other balance measures
Wrisley et al <sup>32</sup>	2007	Reliability of SOT over 2-week period in healthy YA	13	Multiple balance measures needed for more challenging situations, specifically athletes
Yamada et al <sup>33</sup>	2011	Can Wii Fit be used to assess fall risk in OA?	45	Functional measures of health relate significantly to basic step test on Wii Fit

Abbreviations: AT, athletic trainer; BESS, Balance Error Scoring System; COP, center of pressure; CTSIB, Clinical Test of Sensory Organization and Balance; DOF, degrees of freedom; ICC, interclass correlation coefficient; ImpACT, Immediate Post-Concussion Assessment and Cognitive Testing; N/A, not accessible; NCAA, National Collegiate Athletic Association; OA, older adults; PC, postural control; SEBT, Star Excursion Balance Test; SOT, Sensory Organization Test; YA, young adults.



**Table 2. Extended From Previous Page**

Significant Results	Notes	Type of Publication	Rating
Same as synthesis	No evidence given to support use of Wii Fit in concussion management	Book	N/A
Little to no balance assessment with evidence to support use		Conference proceedings	N/A
BESS has moderate to good reliability in most clinical observations	Moderate reliability in populations with concussions	Systematic review	A
Changes in COP oscillations during visual and nonvisual tests reflect cortical changes or alterations in function	Approximate entropy or sample entropy are new normality scales in observations of COP	Original research	A
Same as synthesis		Original research	B
ICC ranged from 0.26 to 0.64	% agreement was 77–100 for the first condition on the SOT	Original research	A
Vestibular is most common interruption after concussion	Clear information regarding current balance techniques	Review of literature	A
Impact location was mainly front and top portion of head	Relationship between magnitude and location influenced return to play	Original research	A
Balance impairment lasted 1 to 3 days postinjury	CTSIB displayed balance impairments 3 days postinjury	Original research	A
Suggest Wii Fit can be used for static balance assessments in Parkinson patients	Parkinson relationship between individuals with concussions could be key	Original research	B
Severely head-injured individuals sway more in Romberg	Background along with practical application of the Romberg	Original research	C
Sensitivity of 0.55 and specificity of 0.64 in YA, and 0.61 and 0.58 respectively for OA; all with vestibular dysfunction diagnosed	Romberg or modified Romberg tests cannot easily detect vestibular problem	Original research	A
Same as synthesis	Detection of positive and negative signs from other neurologic disorders	Review of literature	D
Same as synthesis	Romberg test originally designed for diagnosing tabes dorsalis dysfunction	Review of literature	C
Balance issues persist up to 72 hours after concussion.	Postural-stability testing provides a useful tool for objectively assessing the motor domain of neurofunctioning; considered reliable and valid	Conference proceedings	N/A
Sensitivity = 0.34, specificity = 0.91 to 0.96	Balance deficits dissipated after 3 to 5 days postinjury	Original research	A
Different motor strategies used during Wii Fit games	Supports further use of COP with force plate	Original research	B
Wii Fit does correlate with visual-processing speed	Different motor strategies needed for Wii Fit balance game	Original research	A
Romberg test is inappropriate for measuring balance in athletes with concussions	BESS relates well to SOT	Original research	A
Specific guidelines of CTSIB and how it is conducted	CTSIB is cost-effective way to measure balance	Original research	A
Reliability was ICC = 0.84 with eyes open and 0.86 with eyes closed	Supports reliability of measures of Romberg test	Original research	B
Reliability was ICC = 0.98 in OA	High reliability of CTSIB for use in OA	Review of literature	B
Intrasession ICC range = 0.8–0.39, intersession ICC = 0.74–0.29	Wii Fit balance activity had poor validity relative to COP and SEBT	Original research	A
Reliability mean composite ICC = 0.64	Moderate to low levels of reliability in healthy individuals	Original research	C
Game-based fall risk assessment can be done with Wii Fit basic step test	Interesting relationship between dual tasks and Wii Fit	Original research	B

author (N.M.) made the final judgments regarding inclusion or exclusion of the articles identified.

We found that no reliability or validity data currently support the use of the CTSIB, the SOT, the Romberg test,

or the Wii Fit balance game for evaluating balance dysfunction in concussion injuries. The BESS was the only balance assessment that had supporting reliability and validity evidence for use in this population.<sup>9,11,16,24</sup> In a

single study<sup>18</sup> of reliability, high reliability of the CTSIB was noted in older normal adults, but no reliability or validity evidence was seen in the concussion group (Table 2). The SOT had 2 articles<sup>25,32</sup> supporting moderate to high reliability in healthy young and older adults. No reliability and validity evidence was demonstrated for the SOT in patients with concussions. Two studies<sup>9,11</sup> supported the BESS's moderate to high reliability, and in a single study,<sup>16</sup> high specificity and low sensitivity values were observed. A single study<sup>34</sup> of reliability showed moderate to high reliability (0.86) of the Romberg test for use in diagnosing balance dysfunction in individuals with Parkinson disease.

In the Wii Fit literature search, we found 2 articles<sup>25,31</sup> that supported the use of the Wii Fit for static balance activities and a single article<sup>26</sup> that did not support the use of the Wii Fit in dynamic balance tasks (Table 2). No currently published research studies have examined the validity of the Wii Fit.

## DISCUSSION

The purpose of our systematic review was to examine the reliability and validity evidence for the CTSIB, the SOT, the BESS, the Romberg test, and the Wii Fit balance game for detecting balance disturbance in athletes with concussions. Based on the results of this review, no reliability and validity data currently exist in support of the use of the CTSIB, SOT, Romberg test, or Wii Fit for detecting concussion-related balance disturbances in all demographics. The BESS has high reliability but low validity (specifically sensitivity and specificity values) to assess balance disturbances in athletes with concussions. Because of the limitations in the literature, other neurologic disorders similar to concussion could provide reliable reliability and validity data to support the use of the CTSIB, SOT, Romberg, or Wii Fit tests for detecting balance dysfunction.

### Clinical Test of Sensory Organization and Balance

The CTSIB was originally developed by Shumway-Cook and Horak<sup>28</sup> in 1986 and involves 6 major scenarios that systematically remove or conflict sensory inputs.<sup>6,28</sup> The modified CTSIB, which involves more technically complex equipment measures such as a force platform, is used more commonly in research. The assessment procedures for the CTSIB, regardless of modification, include the following phases: alternating between standing on high-density foam or the ground, with eyes open, closed, or looking into an object.<sup>28</sup> Balance is measured using a scale of 1 to 4: a lower value indicates the least amount of postural sway, and a value of 4 indicates the potential for falling.<sup>28</sup>

Guskiewicz et al<sup>19</sup> administered the modified CTSIB and found that participants with concussions had decreased postural stability compared with baseline scores during the initial 3 days after injury. These results suggest that the balance deficits were related to sensory integration interaction errors. The CTSIB is a basic clinical test and can be administered using clinical measures or laboratory-grade technology, but it is considered a lower standard of assessment.<sup>6</sup> Whitney et al<sup>30</sup> measured the reliability of the unmodified CTSIB in older adults and reported an interclass correlation coefficient (ICC) of 0.98, indicating that the CTSIB is a useful test to assess balance deficits in

**Table 3. Reliability and Validity Measures**

Test	Population	Reliability <sup>a,b</sup>	Sensitivity/ Specificity
Clinical Test of Sensory Organization and Balance	Older adults	0.98	N/A
Sensory Organization Test	Older adults	0.49 <sup>b</sup>	N/A
	Young adults	0.64	N/A
Balance Error Scoring System	Athletes with concussions	0.87 <sup>b</sup>	0.34/0.96
Romberg	Parkinson (eyes open or closed)	0.84/0.86	N/A
	Vestibular impairment (young adults)	N/A	0.55/0.64
	Vestibular impairment (older adults)	N/A	0.61/0.58
	Athletes with concussions	N/A	0.55/0.77
Wii Fit	Healthy young adults (force plate)	0.94	N/A
	Athletes with concussions	N/A	N/A

Abbreviations: COP, center of pressure; N/A, not available.

<sup>a</sup> Refers to the highest reliability rating reported in the studies including interclass correlations, intraclass correlations, and intersession correlations.

<sup>b</sup> ICC values were calculated from range values given in studies.

older adults (Table 3). However, no reliability, sensitivity, or specificity measures have been reported in the literature for healthy young individuals or individuals with concussions. Therefore, generalizability of the CTSIB is limited.

### Sensory Organization Test

The SOT has gained considerable support in recent years and has proved to be a viable measure for assessing balance after concussion.<sup>11</sup> The SOT uses a force platform, a reference point, and a harness cage for safety. The SOT systematically disrupts afferent sensory information by reducing spatial awareness cues via somatosensory or visual components.<sup>11</sup> Individuals must attempt to maintain a steady quiet standing posture during each trial. Six different combinations of 3 trials each that alter the afferent information for 20 seconds provide clinicians with a composite balance score and an equilibrium performance score. Scoring is based on a 100-point scale, and higher scores correlate with better balance. For example, a participant whose eyes are covered will attempt to remain stable as the force platform translates in the anterior-posterior directions. In another combination, the environmental cage is moved in reference to the body's sway without movement of the force platform.

A limited number of authors investigated the reliability of instruments designed to assess balance. For example, the SOT is limited in terms of its reliability. Ford-Smith et al<sup>18</sup> assessed the test-retest reliability of the SOT in 40 noninstitutionalized adults over the age of 65 years. The

ICCs ranged from 0.26 to 0.64 during a week-to-week average of the 3 trials of the SOT balance test (Table 3). This range corresponds to reliability that is highly variable within and across conditions. Wrisley et al<sup>32</sup> assessed the reliability of the SOT over a 2-week period in 15 healthy young adults. The mean composite ICC score was 0.64. The reported differences between the young and older adults could be attributed to age-related deficits in functional stability, age-related changes in neuromuscular strategies to maintain posture, or the differences in multiple testing scenarios. Overall, the SOT shows a moderate level of reliability in healthy individuals, but reliability tests have not been conducted in injured populations; therefore, generalizability is limited.

### Balance Error Scoring System

The BESS was developed at the University of North Carolina–Chapel Hill to provide a low-technology, cost-effective, sport-related concussion assessment for balance.<sup>6</sup> This test requires a foam platform and a stopwatch and can be used as both a sideline and a clinical test. Individuals perform a series of stances that include double, single, and tandem (legs in a heel-to-toe formation) leg support. The starting and reference position, for grading of balance errors, requires the participant to place the hands upon the iliac crest and close the eyes. A 20-second clinical observation period begins once the participant's eyes are closed. Performance is assessed by the number of deviations or errors from the original position, with a maximum of 10 possible errors per trial. Two trials are assessed for each foot position, resulting in 6 total trials.

The BESS is currently the most widely used postconcussion balance test because it is a low-technology form of analysis.<sup>6</sup> In a study of the interrater reliability between the BESS and SOT measurements in athletes with concussions, the ICCs ranged between 0.78 and 0.96.<sup>10</sup> However, when balance was evaluated in individuals with concussions, the sensitivity value for the BESS at baseline was 0.34 and the specificity values range from 0.91 to 0.96 across days 1–7 after the injury (Table 3).<sup>25</sup>

The BESS is generally widely accepted by researchers as an appropriate test for assessing balance deficiencies in individuals with concussions.<sup>15</sup> The BESS has been thoroughly examined and systematically reviewed in many populations, including young adults, old adults, athletes of different sports, and persons with concussions<sup>15</sup>; furthermore, it is emerging as the current gold standard in nonlaboratory settings to evaluate balance deficits.<sup>15</sup> Though the BESS has wide popularity and has been clinically validated, limitations remain. Because of its reported low sensitivity statistic (0.34) and inability to detect balance dysfunction at day 7 after an initial concussion, the BESS may best be used as a prescreening test to augment sideline measurements of a suspected concussion.<sup>6</sup> Lastly, the BESS relies heavily upon rater interpretation, resulting in lower interrater reliability.<sup>15</sup>

### Romberg Test

The Romberg test was initially developed in the late 19th century to detect sensory impairments in individuals with *tabes dorsalis*.<sup>24</sup> The purpose of the Romberg test is to assess balance when individuals experience reduced visual

sensory input. With reduced visual input, reliance on the vestibular and somatosensory systems for balance and postural modulations is exaggerated. Commonly used by physicians and other health care professionals, the Romberg test evaluates potential balance impairments after a neurologic incident such as a concussion.<sup>6,23</sup> Postural control can be examined via the Romberg test by having the individual stand as quietly as possible without deviating, with the feet placed together (left and right medial malleoli touching) or in the heel-to-toe position, with differing visual sensory conditions.<sup>10,11,16</sup> The visual sensory conditions are eyes open and eyes closed.<sup>10,11,16</sup> The Romberg sign is positive if the individual begins to sway involuntarily during the reduced-vision condition.<sup>29</sup> If abnormal movements or sway are detected during the Romberg test, then a neurologic deficit is believed to exist within the higher-functioning sensory systems.<sup>24</sup> It has been suggested that the sensory dysfunction (associated with *tabes dorsalis* or concussion) can be related to myelopathies or neuropathies of the sensory system.<sup>23,24</sup> If myelopathy, neuropathy, or short-term reduction in flow of sensory information to higher centers exists, balance or postural control will be reduced.

Reliability and validity data of the Romberg test are limited. We are aware of no reliability or validity data that have been reported for this test in individuals who have experienced concussions. However, reliability evidence does exist for the Romberg test in other neurologic dysfunctions. The test-retest intraclass correlation for individuals with Parkinson disease was 0.84 in the reduced-vision condition and 0.86 in the normal-vision condition<sup>29</sup> (Table 3). These ICC values quantify the reliability of the Romberg test as excellent in individuals with Parkinson disease.

Therefore, regardless of the similarities in the mechanism underlying postural instability, no reliability or validity data currently exist for the use of the Romberg test in assessing concussion injuries.<sup>7</sup> It is possible that a relationship exists between the sensory and balance deficits observed in a concussion and in Parkinson disease. Specifically, both concussion and Parkinson disease are associated with a reduction in the capacity of the basal ganglia to process sensory input, resulting in insufficient sensory integration and aberrant motor control for postural maintenance.<sup>34</sup>

Over the past years, the Romberg test has evolved because of its subjectivity.<sup>22</sup> The Romberg test of standing balance on firm and compliant support surfaces is more commonly used to identify vestibular dysfunction.<sup>22</sup> The Romberg test of standing balance on firm and compliant support surfaces expands the traditional standing test with the addition of a compliant support surface similar to that used in the BESS.<sup>6,22</sup> The compliant surface emphasizes reliance upon the proprioceptive or vestibular sensory systems, effectively testing the sensory systems most commonly responsible for balance dysfunction.

The sensitivity and specificity values of the Romberg test of standing balance on firm and compliant support surfaces reported in individuals with vestibular dysfunction were 0.55 and 0.64, respectively. In individuals over the age of 40 years, the sensitivity and specificity values were 0.61 and 0.58, respectively<sup>22</sup> (Table 3). Thus, the Romberg test accurately detects balance dysfunction in approximately



60% of cases and should not be used as a screening measure for vestibular impairment.<sup>22</sup>

## Wii Fit

The Wii Fit balance game uses visual display to transpose lower body ground reaction force data to meet certain goals based upon the game's requirements. The Wii Fit has been demonstrated to be a reliable measure for static standing balance assessments of center of pressure when compared with the gold standard laboratory-grade force platform.<sup>17</sup> The ICC was 0.94 for results within the device and 0.89 between the Wii Fit and a laboratory-grade force platform (Table 3).<sup>17</sup> Furthermore, the Wii Fit balance game improves anterior-posterior sway during quiet stance in older adults when used as a treatment instrument.<sup>17</sup>

Examining the use of the Wii Fit in patients with Parkinson disease could provide additional insight into reliability and validity data for those with concussions. In a comparable static standing condition, center-of-pressure assessments revealed ICCs ranging between 0.92 and 0.98 when a force platform was compared with the Wii Fit (Table 3).<sup>20</sup> This evidence supports the suggestion that the Wii Fit can be used as an effective measure of static balance and is comparable with the gold-standard force platform.<sup>17,20</sup> However, the Wii Fit may be less accurate during active balance tasks that involve sudden dynamic movements.

During active movements of participants with lower extremity injury, the Wii Fit had poor concurrent validity relative to center of pressure and Star Excursion Balance Test measurements.<sup>31</sup> The intersession correlation coefficient ranged from 0.39 to 0.80, and the intrasession reliability scores ranged from 0.29 to 0.74.<sup>31</sup> Given the large ranges of reliability scores during active movements, the Wii Fit appears to exhibit poor reliability and validity in individuals with lower extremity injuries and in dynamic movements.

The Wii Fit is currently being used by fitness and rehabilitation clinicians to assess and aid in the recovery from injury. The range of Wii Fit games challenges a variety of motor-control strategies to accomplish the predetermined goals of the game.<sup>26</sup> Therefore, it is essential for researchers to identify which type of Wii Fit game appropriately assesses balance. In community-dwelling older adults, dual-task walking and dual-task timed up-and-go maneuvers have been significantly correlated with the basic Wii Fit step test.<sup>33</sup> The Wii Fit basic step test is a replication of the basic aerobic step test, which is considered a functional health assessment for submaximal and maximal cardiorespiratory fitness.<sup>35</sup> However, the validity and reliability measures of balance assessment in individuals with concussions have not been established for the Wii Fit. Additionally, no researchers have demonstrated a clear relationship between the Wii Fit and the other standard clinical assessments of balance or commonly used postural-control assessments in concussion management.

## CONCLUSIONS

The findings of our systematic review suggest that no published reliability or validity data exist for the CTSIB, the SOT, the Romberg test, or Wii Fit in assessing balance

in athletes with concussions. The BESS has high reliability and low validity evidence (specifically sensitivity) to support its use in assessing balance in an athlete with a concussion. The Romberg test and Wii Fit are reliable tools in Parkinson patients. Further reliability and validity data are needed to determine if the CTSIB, SOT, Romberg test, and Wii Fit can accurately assess balance in athletes with concussions. Additionally, more studies that evaluate the relationship between current clinical or standard balance assessments and the Wii Fit are needed.

## REFERENCES

1. McCrea M, Hammeke T, Olsen G, Leo P, Guskiewicz K. Unreported concussion in high school football players: implications for prevention. *Clin J Sport Med*. 2004;14(1):13–17.
2. Gessel LM, Fields SK, Collins CL, Dick RW, Comstock RD. Concussions among United States high school and collegiate athletes. *J Athl Train*. 2007;42(4):495–503.
3. Aubry M, Cantu R, Dvorak J, et al. Summary and agreement statement of the First International Conference on Concussion in Sport, Vienna 2001: recommendations for the improvement of safety and health of athletes who may suffer concussive injuries. *Br J Sports Med*. 2002;36(1):6–10.
4. McCrory P, Johnston K, Meeuwisse W, et al. Summary and agreement statement of the 2nd International Conference on Concussion in Sport, Prague 2004. *Br J Sports Med*. 2005;39(4):196–204.
5. McCrory P, Meeuwisse W, Aubry M, et al. Consensus statement on concussion in sport: the 4th International Conference on Concussion in Sport held in Zurich, November 2012. *J Sci Med Sport*. 2013;16(3):178–189.
6. Guskiewicz KM. Balance assessment in the management of sport-related concussion. *Clin Sports Med*. 2011;30(1):89–102, ix.
7. Guskiewicz KM, Mihalik JP, Shankar V, et al. Measurement of head impacts in collegiate football players: relationship between head impact biomechanics and acute clinical outcome after concussion. *Neurosurgery*. 2007;61(6):1244–1252.
8. Marar M, McIlvain NM, Fields SK, Comstock RD. Epidemiology of concussions among United States high school athletes in 20 sports. *Am J Sports Med*. 2012;40(4):747–755.
9. Slobounov S, Slobounov E, Sebastianelli W, Cao C, Newell K. Differential rate of recovery in athletes after first and second concussion episodes. *Neurosurgery*. 2007;61(2):338–344.
10. Riemann BL, Guskiewicz KM. Effects of mild head injury on postural stability as measured through clinical balance testing. *J Athl Train*. 2000;35(1):19–25.
11. Guskiewicz KM, Ross SE, Marshall SW. Postural stability and neuropsychological deficits after concussion in collegiate athletes. *J Athl Train*. 2001;36(3):263–273.
12. NCAA Publications. *NCAA Sports Medicine Handbook*. Updated 2010–2011. Indianapolis, IN: National Collegiate Athletic Association; 2010.
13. Looney MA. Measurement issues in the clinical setting. In: Wood TM, Zhu W, eds. *Measurement Theory and Practice in Kinesiology*. Champaign, IL: Human Kinetics; 2006:297–303.
14. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol*. 2009;62(10):e1–e34.
15. Bell DR, Guskiewicz KM, Clark MA, Padua DA. Systematic review of the Balance Error Scoring System. *Sports Health*. 2011;3(3):287–295.
16. Cavanaugh JT, Guskiewicz KM, Stergiou N. A nonlinear dynamic approach for evaluating postural control: new directions for the

- management of sport-related cerebral concussion. *Sports Med (Auckland NZ)*. 2005;31(11):935–950.
17. Clark RA, Bryant AL, Pua Y, McCrory P, Bennell K, Hunt M. Validity and reliability of the Nintendo Wii Balance Board for assessment of standing balance. *Gait Posture*. 2010;31(3):307–310.
  18. Ford-Smith CD, Wyman JF, Elswick RK Jr, Fernandez T, Newton RA. Test-retest reliability of the sensory organization test in noninstitutionalized older adults. *Arch Phys Med Rehabil*. 1995;76(1):77–81.
  19. Guskiewicz KM, Perrin DH, Gansneder BM. Effect of mild head injury on postural stability in athletes. *J Athl Train*. 1996;31(4):300–306.
  20. Holmes JD, Jenkins ME, Johnson AM, Hunt MA, Clark RA. Validity of the Nintendo Wii® balance board for the assessment of standing balance in Parkinson's disease. *Clin Rehabil*. 2013;27(4):361–366.
  21. Ingersoll CD, Armstrong CW. The effects of closed-head injury on postural sway. *Med Sci Sports Exerc*. 1992;24(7):739–743.
  22. Jacobson GP, McCaslin DL, Piker EG, Gruenwald J, Grantham S, Tegel L. Insensitivity of the “Romberg test of standing balance on firm and compliant support surfaces” to the results of caloric and VEMP tests. *Ear Hear*. 2011;32(6):e1–e5.
  23. Khasnis A, Gokula RM. Romberg's test. *J Postgrad Med*. 2003;49(2):169–172.
  24. Lanska DJ, Goetz CG. Romberg's sign: development, adoption, and adaptation in the 19th century. *Neurology*. 2000;55(8):1201–1206.
  25. McCrea M, Guskiewicz KM, Marshall SW, et al. Acute effects and recovery time following concussion in collegiate football players: the NCAA Concussion Study. *JAMA*. 2003;290(19):2556–2563.
  26. Michalski A, Glazebrook CM, Martin AJ, et al. Assessment of the postural control strategies used to play two Wii Fit videogames. *Gait Posture*. 2012;36(3):449–453.
  27. Reed-Jones RJ, Dorgo S, Hitchings MK, Bader JO. Wii Fit plus balance test scores for the assessment of balance and mobility in older adults. *Gait Posture*. 2012;36(3):430–433.
  28. Shumway-Cook A, Horak FB. Assessing the influence of sensory interaction of balance: suggestion from the field. *Phys Ther*. 1986;66(10):1548–1550.
  29. Steffen T, Seney M. Test-retest reliability and minimal detectable change on balance and ambulation tests, the 36-item short-form health survey, and the unified Parkinson disease rating scale in people with parkinsonism. *Phys Ther*. 2008;88(6):733–746.
  30. Whitney SL, Poole JL, Cass SP. A review of balance instruments for older adults. *Am J Occup Ther*. 1998;52(8):666–671.
  31. Wikstrom EA. Validity and reliability of Nintendo Wii Fit balance scores. *J Athl Train*. 2012;47(3):306–313.
  32. Wrisley DM, Stephens MJ, Mosley S, Wojnowski A, Duffy J, Burkard R. Learning effects of repetitive administrations of the sensory organization test in healthy young adults. *Arch Phys Med Rehabil*. 2007;88(8):1049–1054.
  33. Yamada M, Aoyama T, Nakamura M, et al. The reliability and preliminary validity of game-based fall risk assessment in community-dwelling older adults. *Geriatr Nurs*. 2011;32(3):188–194.
  34. Kandel ER, Schwartz JH, Jessel TM. *Principles of Neural Science*. 4th ed. New York, NY: McGraw-Hill; 2000:810–814.
  35. Howley ET, Thompson DL. *Fitness Professional's Handbook*. 6th ed. Champaign, IL: Human Kinetics; 2012.

---

Address correspondence to Nicholas Murray, PhD, Department of Health and Kinesiology, Georgia Southern University, PO Box 8076, Statesboro, GA 30460. Address e-mail to nmurray@georgiasouthern.edu.