ACL Research Retreat VII: An Update on Anterior Cruciate Ligament Injury Risk Factor Identification, Screening, and Prevention

March 19-21, 2015; Greensboro, NC

Sandra J. Shultz, PhD, ATC, FNATA, FACSM*; Randy J. Schmitz, PhD, ATC*; Anne Benjaminse, MSc, PT†; Malcolm Collins, PhD‡; Kevin Ford, PhD§; Anthony S. Kulas, PhD, ATCII

*Department of Kinesiology, University of North Carolina at Greensboro; †The Department of Human Movement Sciences, University of Groningen, and The School of Sports Studies, Hanze University, Groningen, The Netherlands; ‡Department of Human Biology, University of Cape Town, South Africa; §Human Biomechanics and Physiology Laboratory, Department of Physical Therapy, High Point University, NC; IIDepartment of Health Education and Promotion, Eastern Carolina University, Greenville, NC

he seventh ACL Research Retreat was held March 19–21, 2015, in Greensboro, North Carolina. The retreat brought together clinicians and researchers to present and discuss the most recent advances in anterior cruciate ligament (ACL) injury epidemiology, risk factor identification, and injury risk screening and prevention strategies. Subsequently, our goal was to identify important unknowns and future research directions.

The ACL Research Retreat VII was attended by 64 clinicians and researchers from Australia, Canada, India, Ireland, the Netherlands, South Africa, the United States, and the United Kingdom. The meeting featured 3 keynote and 29 podium presentations highlighting recent research. Keynotes were delivered by Bruce Beynnon, PhD (University of Vermont), Charles "Buz" Swanik, PhD, ATC (University of Delaware), and Mark Paterno, PhD, PT, ATC, SCS (Cincinnati Children's Hospital Medical Center), addressing their ongoing work related to sex-specific multivariate risk factor models for ACL injury,¹ the role of the brain in noncontact ACL injury,² and the incidence and predictors of a second ACL injury after primary ACL reconstruction and return to sport,³ respectively. Podium and poster presentations were organized into thematic sessions of prospective and case-control risk factor studies, anatomical and hormonal risk factors, neuromuscular and biomechanical risk factors, injury risk assessment after ACL injury, and injury-prevention strategies. Time was provided for group discussion throughout the conference. At the end of the meeting, attendees participated in 1 of 3 breakout sessions on the topics of genetic, hormonal, and anatomical risk factors; neuromechanical contributions to ACL injury; and risk factor screening and prevention. From these discussions, we updated the 2012 consensus statement⁴ to reflect the most recent advances in the field and to

revise the important unknowns and future directions necessary to enhance our understanding of ACL injury. Following are the updated consensus statement, keynote presentation summaries, and free communication abstracts organized by topic and presentation order.

CONSENSUS STATEMENT

As our understanding of ACL injury risk and prevention grows, it is becoming increasingly apparent that ACL injury mechanism(s) is(are) multifactorial, resulting from the interplay of neuromuscular, biomechanical, anatomical, genetic, hormonal, and other factors. Although we have learned much in recent years about sex differences in neuromechanical movement patterns and the external loading factors that strain the ACL, far less is known about the factors that contribute to a structurally weaker ligament. This is important because the effect of external loading factors on injury risk potential (ie, the potential to cause ligament failure) likely depends on the intrinsic properties (ie, structural integrity and load-bearing capabilities) of the ligament. From this perspective, new evidence continues to emerge regarding the potential for one's genetic makeup and hormonal profile to substantially alter the structural properties of the ACL, making it more or less vulnerable to failure with external loading. This interplay between intrinsic ligament properties and external loading may explain why screening movement patterns alone has yet to yield consistent and reliable risk factor prediction models. Additionally, consensus is building that the risk factors may not be the same for males and females. As such, examining this interplay would help us to further elucidate whether females are at greater risk of noncontact ACL injury due to female-specific injury mechanisms or the same external-loading injury mechanisms

apply to both sexes, but the underlying intrinsic risk factors are merely more prevalent in females.

Despite the likely multifactorial nature of this injury, studies of isolated risk factors (eg, anatomic, hormonal, genetic, kinematic, or kinetic) continue to dominate the literature. Further, because of the relatively low incidence of ACL injury, lower extremity biomechanics are often studied as outcome measures to represent ACL injury potential based on what we have learned from cadaveric ligament-loading studies. If we are to fully understand the movement behavior that results in elevated ACL loading, researchers need to study ACL injury as the outcome and move toward a more comprehensive assessment of both the nonmodifiable (ie, anatomy, genetics, and hormones) and modifiable (ie, neuromechanics) factors that influence risk. Conducting such multifactorial studies with injury as the outcome will allow us to determine important interactions and interdependencies among the various risk factors and identify the unique combination of factors that can most reliably predict future injury risk potential in the simplest way possible. Although tremendous challenges remain in performing such large-scale multivariate risk factor studies (eg, funding, personnel, and geographic restrictions), results from these types of studies are beginning to emerge and published findings have been incorporated into the current statement where available. Additional and important insights are also being gained from the studies of the incidence and predictors of a second ACL injury after primary ACL reconstruction and full return to sport participation; this information has also been added to the statement (see also Paterno³).

Because of the trends noted earlier, this consensus statement differs somewhat in format compared with previous statements. The first section focuses primarily on risk factor assessment as determined by ACL injury as the outcome variable. However, we have also included biomechanical studies that describe how the ACL is stressed or strained in order to better understand mechanistically how global movements (and the intrinsic and extrinsic factors that affect them) may lead to ACL failure. Although this section is still divided into the traditional risk factor categories, we have made every effort to integrate the published findings from multivariate studies as they became available. Also new to this section is information on the risk of second-time ACL injury. The second and third sections focus on risk factor screening and prevention efforts, respectively.

Once again, we find that in the 3 years since the last ACL Research Retreat, many advances in our knowledge have shaped what we know about ACL injury risk and prevention and the important directions for future research that are needed to move the field forward. We hope that these proceedings will continue to stimulate quality research to more effectively identify those at risk before an injury occurs and to promote high-quality clinical interventions.

RISK FACTOR ASSESSMENT

Neuromechanical Risk Factors

Neuromechanical (ie, neuromuscular and biomechanical) factors, whether ascertained in vivo or in vitro, are

generally derived from instrumented analyses of function that typically include kinematics, kinetics, and the timing and magnitude of muscular activation and force production. Our understanding of the role of neuromechanical factors in ACL injury has been advanced through cadaveric modeling, movement biomechanics, and neuromuscular insufficiency.

Important Knowns and Recent Advances. Biomechanics Influencing ACL Strain. Advances in our understanding of how the ACL is loaded and how it fails have arisen from cadaveric and computer models of simulated knee loads. These studies are foundational if we are to better understand and interpret "high-risk" movement biomechanics studies that do not estimate ACL loads or use ACL injury as an outcome. The ACL is loaded in vitro through a variety of isolated and combined mechanisms that are assumed to be present during dynamic sport postures considered to be high risk. Collectively, this work demonstrates that ACL strain increases with anterior tibial forces, tibiofemoral compression, and combined knee-abduction and kneeinternal-rotation moments.^{5–13} Mechanical couplings of external moments about the knee that include combined internal rotation and abduction are reported to increase ACL strains to near levels at which tissue ruptures.⁸

Recent cadaveric studies have applied loads to the knee that simulate activities representing high-risk ACL injury events. The timing of peak ACL strain appears to depend more on anteriorly directed tibial forces and knee-abduction moments, with peak internal tibial rotation occurring much later in the simulated landing.¹⁰ In addition, knee abduction combined with internal knee rotation, anterior tibial translation, and increased tibial compression produces ACL injuries that appear consistent with clinical observations of ACL injuries.¹⁴ During simulated single-legged landing, peak ACL strain increases with a combination of larger ground reaction forces and decreased hip angles.¹⁵ Collectively, these cadaveric studies have clarified how the ACL is loaded and injured and formed the basis for assessing biomechanical variables thought to represent "high-risk" movement biomechanics.

Movement Biomechanics. Noncontact ACL injuries are most commonly reported to arise from a sudden deceleration while changing direction when running or landing from a jump.¹⁶ Quantitative analyses of actual injury events demonstrate that rapid knee abduction and internal rotation during the early weight-bearing phase occur at the time of injury.^{17,18} Additionally, video observations of ACL injury occurrences demonstrate a relatively extended knee,¹⁹ greater knee-abduction motion,^{20–22} increased amounts of lateral trunk motion,²³ and a more posteriorly positioned center of mass.²⁴ These observational findings are largely supported by cadaveric model work and lend credence to the use of high-risk biomechanical measures as outcome variables.

To date, 2 groups have demonstrated landing biomechanics to be prospective risk factors of ACL injury. When researchers used full 3-dimensional biomechanical methods, larger peak knee-abduction moment and peak kneeabduction angle at initial contact during a drop jump were prospective risk factors in adolescent female athletes.²⁵ Use of a more clinically friendly video observation of a jump landing (Landing Error Scoring System [LESS]) in youth soccer athletes showed that participants who went on to sustain ACL tears had lower LESS scores.²⁶ However, LESS scores did not predict ACL injury in high school and college-aged athletes.²⁷

A more erect or upright posture is commonly associated with increased vertical ground reaction forces.^{28,29} This is important to understand as in vivo and in vitro strain of the ACL is related to maximal load and timing of ground reaction forces.^{30,31} Similarly, anterior tibial translation increases as demands on the quadriceps increase.^{32,33} Thus, this upright or extended posture when contacting the ground during the early stages of deceleration tasks has been suggested to be associated with the ACL injury mechanism.^{34–38} Although short-term exhaustive exercise has resulted in changes in non–sagittal-plane variables associated with ACL injury risk,^{39–43} exertional protocols that aim to simulate entire game or match duration and intensity have demonstrated subtle changes only in the sagittal plane, indicative of an at-risk upright posture.^{44,45}

Females are known to be at greater risk of ACL injury than males,^{46,47} so investigations have focused predominantly on sex differences. Several recent reviews^{48,49} have indicated that females demonstrated greater knee-abduction postures during landing. Given the prospective findings of Hewett et al,²⁵ knee abduction may be a potential key biomechanical factor in defining female-specific ACL injury mechanisms.⁴⁹

Neuromuscular and Neurocognitive Insufficiency. Trunk neuromuscular control has been implicated as a risk factor for ACL injury. In a prospective study⁵⁰ of trunk displacement after sudden force release, greater trunk displacement was a prospective predictor of ACL injury in college-aged females but not males. Compared with an erect or preferred trunk posture, landing with the trunk relatively more flexed increased hip-flexion and hip-extensor moments while decreasing ground reaction forces, knee moments, and quadriceps activation amplitudes.^{28,51} During various running and cutting tasks, trunk and hip movements (in all 3 planes of motion) were commonly associated, at different magnitudes, with knee-abduction angles and external kneeabduction moments.^{52–55} Although trunk and hip associations with "high-risk" knee biomechanics are clearly supported by these latter studies, the correlational results do not suggest cause and effect and could instead reflect task-dependent associations. Collectively, positions and movements of segments proximal to the knee could place the knee joint in a position of high risk for ACL injury.

Neurocognitive insufficiency as a risk factor for ACL injury is supported by findings of worse baseline reaction times, processing speed, and visual-spatial awareness in athletes who went on to sustain an ACL injury compared with controls.^{2,56} Evidence for a connection between central processing and neuromuscular insufficiency was provided by the finding that slower cognitive processing speed was associated with decreased trunk stability in males.⁵⁷

Unknowns and Directions for Future Research.

1. External- and internal-loading profiles that cause noncontact ACL ruptures in the field, data that are central to optimal injury-prevention strategies, are unknown.⁵⁸ Video from actual injury situations (and control videos from these athletes before injury) must be accumulated^{17,18,24} to allow us to better understand the mechanisms of in vivo ACL rupture. Additionally, cadaveric, mathematical, in vivo kinematic, and imaging research approaches should be combined to help us understand the postures, loads, and neuromuscular profiles that increase ACL strain or cause noncontact ACL rupture.^{31,59}

- 2. Optimal ways to assess movement in the laboratory environment are still being debated with regard to single-legged and double-legged tasks.^{60,61} To better understand how movement patterns and other structures in the kinetic chain affect ACL loads, we must continue to develop, improve, and validate quality laboratory-based models (eg, computational, cadaveric) that noninvasively estimate in vivo ACL forces and strain. Care should be taken to not overgeneralize results from 1 specific task to other tasks.
- 3. The influence of the kinetic chain (ie, ankle, hip, lumbopelvic, and trunk biomechanics) on modeled ACL strain or forces has been studied in controlled movements commonly performed in rehabilitation.^{62,63} However, the specific relationship of these kinetic chain concepts with ACL injury risk is unclear. Laboratory, cadaver, and simulation-based studies designed to evaluate cause and effect (ie, highly controlled human movement studies with 1 variable manipulated) are warranted to ascertain whether these kinetic chain factors are a cause of or compensation for poor knee biomechanics.
- 4. Typical biomechanical experiments rely on extracting singular measures to characterize complex human biomechanics. To better understand the richness of these data, we should consider alternative approaches to analyzing traditional biomechanical data: for example, nonlinear dynamics⁶⁴ and emerging data-analysis techniques, including statistical parametric mapping.^{65–67}
- 5. To better understand the role of the central nervous system in ACL injury, research models should include assessments of central processes (eg, automaticity, reaction time), cognitive processes⁶⁸ (eg, decision making, focus and attention, prior experience [expert versus novice]), and metacognitive processes (eg, monitoring psychomotor processes). The reader is referred to Swanik,² who briefly reviewed anecdotal, theoretical, and recent clinical research evidence attempting to unravel the brain's role in maintaining joint stability.
- 6. Although investigation of an initial cadaveric model showed that the ACL is susceptible to fatigue failure during repetitive simulated pivot landings and a smaller cross-sectional area of the ACL is at greater risk for such failure,⁶⁹ we do not know if it is a single episode or multiple episodes (or both) that causes gross failure of the ACL.
- 7. To various degrees, the amount of maximal strength is related to movement mechanics, but how this is related to injury risk is unclear.^{70–75} Further work on how neuro-muscular properties beyond absolute strength (eg, muscle and joint stiffness, muscle mass, rate of force production) relate to ACL function and injury risk is warranted.
- 8. The general consensus is that maturation influences biomechanical and neuromuscular factors affecting ACL strain.^{76–87} However, evidence for how maturation affects knee biomechanics is inconsistent.⁸⁸ Examining the influence of the maturational process on knee biomechanics and specifically on ACL loads may provide unique insights into the observed difference in injury rates by sex that emerges during the early stages of physical maturation. In this regard, multiyear, longitudinal studies are needed.

9. Methodologic concerns regarding data collection, processing, and analysis continue to confound the interpretation of biomechanical variables thought to be risk factors for ACL injury. Concerns related to validity³¹ and reliability^{89–91} need further study before we can trust the quality and consistency of the data. Data processing should be carefully considered for its potential effect on identifying high-risk individuals and associations with ACL injury risk.^{92,93} The research community would also benefit from operational definitions of terms (eg, stiffness, stability) so that scientists and clinicians can more transparently interpret the results and conclusions.

Anatomical and Structural Risk Factors

The primary anatomical and structural factors examined with regard to ACL injury risk include knee-joint geometry, knee-joint laxity, body composition, and lower extremity structural alignments. Most of the variables that account for these anatomical characteristics differ by sex, which suggests that the risk factors may not be the same for males and females. Although static alignment measurements have yet to emerge as strong predictors of ACL injury, evidence associating knee-joint geometry, kneejoint laxity, and body mass index with injury risk continues to build.

Important Knowns and Recent Advances. *Knee-Joint Geometry.* The majority of prospective and retrospective case-control studies comparing ACL-injured patients with uninjured controls indicate that ACL-injured patients have smaller ACLs (in area and volume),^{94,95} greater lateral posterior-inferior tibial plateau slopes (but not necessarily medial tibial slopes),^{96–100} and smaller femoral notch widths and notch width indexes.^{95,98,101–108} The presence of a more prominent or thicker bony ridge on the anteromedial outlet of the femoral intercondylar notch has also been reported in ACL-injured patients versus controls.^{95–98} In addition, reduced condylar depth of the medial tibial plateau may be characteristic of ACL injury.^{96,97,100} Coronal slope has not been associated with ACL injury risk.^{97,100}

Multivariate analyses suggest that smaller ACL volume and femoral notch width at the inlet and outlet and greater anteromedial bony ridge thickness and posterior-inferior directed slope of the lateral tibial compartment may independently contribute to injury risk and together provide more information on injury risk potential.^{95,98,109,110} However, because the geometric variables examined have differed, there is no clear consensus as to the unique combination of these variables that most strongly predicts ACL injury risk in males and females.

When compared with males, females have smaller ACLs as assessed by length, cross-sectional area, and volume, even after adjusting for body size.¹¹¹ After adjusting for age and anthropometric factors, the female ACL has less collagen fiber density (area of collagen fibers/total area of the micrograph)¹¹² and decreased mechanical properties, including strain at failure, stress at failure, and modulus of elasticity.¹¹³ Females are also reported to have greater lateral and medial posterior-inferior tibial slopes,^{114,115} reduced coronal tibial slopes,¹¹⁴ taller femoral notch heights, and smaller femoral notch widths than males.¹¹¹ Femoral notch width and angle are reported to be good predictors of ACL size (area and volume) in males but not

in females¹¹¹; multivariate analyses suggest that the individual characteristics may provide unique information¹¹⁰ and collectively influence femoral notch impingement theory.¹¹¹ Sex differences in knee structural anatomy may also explain why associations of these variables with ACL injury history are somewhat different between males and females.^{95,98,110}

Biomechanically, good evidence indicates that knee-joint geometry is related to higher-risk biomechanics. Greater posterior-inferior lateral tibial slopes are associated with greater anterior joint reaction forces,¹¹⁶ anterior translation of the tibia relative to the femur,^{117,118} and peak anterior tibial acceleration¹¹⁹; when combined with a smaller ACL cross-sectional area, these factors are associated with greater peak ACL strains.¹²⁰ Greater relative posterior-inferior slope of the lateral versus the medial tibial plateau has been associated with greater peak knee-abduction and internal-rotation angles,^{116,121} whereas a reduced coronal slope has been associated with greater hip adduction and knee valgus upon landing in females.¹²¹

Knee-Joint Laxity. Greater magnitudes of anterior knee laxity,^{122,123} genu recurvatum,^{122,124–127} general joint laxity,^{84,122,124,127} and internal-rotation knee laxity¹²⁸ have been reported in the contralateral knees of ACL-injured patients compared with controls.

Knee-joint laxity varies considerably among individuals, likely because of a combination of genetic,¹²⁹ hormonal,^{130–135} and anatomical^{136–138} factors. On average, females have greater sagittal (anterior knee laxity, genu recurvatum)-,^{104,122,139–143} frontal (varus-valgus rotation)-, and transverse (internal-external rotation)-plane^{135,144–146} and generalized joint laxity compared with males.^{104,122} Sex differences in frontal-plane and transverse-plane knee laxity persist, even in males and females with similar sagittal-plane knee laxity.^{135,144,146} Females are also more likely to experience acute increases in knee laxity during exercise¹⁴⁷ and across the menstrual cycle.^{130–135}

Greater magnitudes of knee laxity may have both biological and biomechanical consequences. Evidence from animal studies^{148–150} suggests that greater knee laxity is associated with ligament biomarkers indicative of greater collagen turnover, more immature cross-links, and lower failure loads. Biomechanically, greater magnitudes of knee laxity and general joint laxity have been associated with higher-risk landing strategies more often observed in females (eg, greater knee-extensor loading and stiffening, greater dynamic knee valgus), particularly in the planes of motion where greater knee laxity is observed.^{139,151–157} The acute changes in knee laxity observed during exercise and across the menstrual cycle are of sufficient magnitude to push an individual toward higher-risk movement strategies.^{155,156,158–160}

Lower Extremity Alignment. Although the results of 2 large prospective studies are not yet available¹⁶¹ (see also Beynnon et al¹), no clear consensus in the literature consistently links any 1 lower extremity alignment factor to ACL injury. Lower extremity alignment differs among maturational groups and also develops at different rates in males and females within those groups.¹⁶² Once fully matured, females have greater anterior pelvic tilt, hip anteversion, tibiofemoral angle, and quadriceps angle.^{142,163} No sex differences have been observed in measures of tibial torsion,¹⁴² navicular drop,^{142,143,163} or rearfoot angle.^{142,164}

Body Composition. Increasing evidence from 2 large, prospective, multivariate risk factor studies showed that an elevated body mass index predicts future ACL injury in females but not in males.¹⁰⁴ (See also Beynnon et al.¹) Higher body mass index in females is more likely to reflect a higher proportion of fat mass versus lean mass relative to body weight^{165,166}; less relative lean mass has been associated with greater knee-joint laxity¹³⁷ and dangerous biomechanical strategies.^{51,167} This may explain why the combination of greater joint laxity and greater body mass index is a stronger predictor of ACL injury risk in females than either variable alone.¹⁰⁴ (See also Beynnon et al.¹)

Unknowns and Directions for Future Research.

- 1. Anatomical and structural factors have often been examined independently or in small subsets of variables. Even among 3 larger, prospective multivariate risk factor studies^{104,161} (see also Beynnon et al¹), the anatomical variables examined are quite disparate, making it difficult to build a clear consensus on the best prediction model of ACL injury risk for screening purposes. Further large-scale risk factor studies that account for all relevant lower extremity anatomical and structural factors are needed to reach this consensus. Prospective confirmatory analyses of currently identified risk factor models across different races and ages are also needed.
- 2. Most investigations of tibial plateau geometry are based on measures of the subchondral bone, and recent research¹¹⁰ suggests that accounting for the overlying cartilage and meniscal geometry when characterizing plateau slope and depth may improve prediction. However, because these measures are complex and costly, it will be important to determine the extent to which they provide additional information on ACL injury risk once more clinically accessible risk factors are taken into account. (See also Beynnon et al.¹)
- 3. Studies examining the combined effects of joint laxity, tibial geometry (lateral tibial slope, medial:lateral tibial slope ratio, coronal slope, medial condylar depth) and ACL morphology, as well as interactions among these variables, on tibiofemoral joint biomechanics and ACL strain and failure are encouraged.
- 4. Although anatomical and structural factors are often thought to be nonmodifiable, we have limited knowledge of how these structural factors change during maturation or whether physical activity (or other chronic external loads) can influence this development over time, particularly during the critical growth periods. Prospective longitudinal studies are needed to help us understand the underlying factors that result in at-risk anatomical and structural profiles during maturation while also considering relevant modifiable factors such as body composition, neuromuscular properties, and physical activity.
- 5. Recent evidence¹⁶⁸ suggests that individuals with specific laxity profiles may perceive functional deficits during activities of daily living and sport. We do not yet know the multiplanar laxity profiles that pose the greatest risk for ACL injury or the threshold at which the magnitude of knee laxity becomes problematic.
- 6. Because body mass index is not a true measure of body composition or body mass distribution, future studies using a more detailed analysis of body composition are needed to fully understand how body composition contributes to faulty lower extremity biomechanics and injury risk.

Genetic Risk Factors

Rupture of the ACL is a multifactorial condition. Increasing evidence from both familial and case-control genetic association studies indicates that genetic sequence variants play an important role in its occurrence.

Important Knowns and Recent Advances. Since the previous ACL retreat, a growing number of common DNA sequence variants within genes involved in various biological processes have been associated with susceptibility to ACL rupture. These include variants within several genes that encode collagens^{169–175} and proteoglycans¹⁷⁶ involved in the formation of the collagen fibril, the basic building block of ligaments. A subset of these variants has been specifically associated with ACL ruptures in females but not in males.^{171,173,174,177} The ACL's collagen fibril and the rest of its extracellular matrix continuously undergo remodeling after mechanical loading to maintain tissue homeostasis. In support of this, variants within genes that encode proteins involved in cell-signaling pathways, such as angiogenesisassociated signaling¹⁷⁸ and the apoptosis-signaling cascade,¹⁷⁹ as well as remodeling (specifically matrix metalloproteinases^{180,181}), have also been associated with ACL ruptures. Many of the anatomical, structural, and other risk factors are themselves multifactorial phenotypes determined by, to a lesser or greater extent, both genetic and environmental factors.¹⁸² Based on the current evidence, it is therefore unlikely that the identified genetic variants are independent risk factors and more likely that they modulate risk through their effects on structural differences and other biological variations.

Genetic association studies in humans have recently been strengthened by a publication reporting the association of variants within genes, including collagen and other genes proposed to have a detrimental effect on ligament structure and strength, with canine cranial cruciate ligament rupture.¹⁸³ Using a genome-wide association study design, the same investigators¹⁸⁴ have also identified 3 main chromosomal regions, which include variants within neurologic pathway genes, associated with canine cranial cruciate ligament rupture.

Unknowns and Directions for Future Research.

- 1. Most of the genetic risk factors for ACL rupture to date have been identified in white populations, and the reported associations cannot necessarily be extrapolated to other population groups. For example, the functional Sp1binding site polymorphism (rs1800012, G/T) within the *COL1A1* gene is associated with ACL rupture in white populations.^{169,170,172,175} However, the frequency of the minor T allele of this variant is much lower in many other population groups (www.ensembl.org) and therefore unlikely to be informative within all populations.¹⁸⁵ Hence, investigators should identify appropriate informative genetic markers for other population groups.
- 2. Although establishment of an international consortium or registry was recommended at the last ACL retreat,¹⁷¹ most of the published case-control genetic association studies have involved relatively small sample sizes, especially with respect to the sex-specific genetic effects. The establishment of an international consortium or registry will allow us to test the hypothesis of a stronger genetic contribution in individuals who tear their contralateral ACL rather than retear the same ACL or incur no further

ACL injury. The establishment of an international consortium will also make it possible to recruit large sample sizes for whole-genome screening methods, enabling us to identify all the potentially important biological pathways involved in ACL ruptures and how these might differ by population.

- 3. None of the associated variants alone cause ACL ruptures. Rather, the injury is the result, at least in part, of a poorly understood, complex interaction of external loading and other factors, as well as intrinsic stimuli, with the genetic background of the individual. Further research is needed to ascertain the extent to which the associated variants may result in interindividual variations in the structure and, by implication, the mechanical properties of the ACL and surrounding tissues, as well as its responses to mechanical loading and other stimuli, thereby affecting the risk of injury.
- 4. Although scientists are only starting to understand the contribution of genetics to ACL injury, numerous companies are marketing direct-to-consumer genetic tests for common injuries, including ACL ruptures. However, these tests are premature because the genetic data are incomplete and have not been considered together with clinical indicators and lifestyle factors, which may allow an appropriately qualified health care professional to identify an altered risk for injury.

Hormonal Risk Factors

Sex hormones likely underlie many of the sex-specific characteristics that emerge during puberty. In particular, the large magnitudes of and monthly variations in estrogen and progesterone concentrations that females experience continue to be active areas of ACL injury risk factor research. The hormone relaxin has also gained attention in recent years for its potential to influence ACL structural integrity and injury risk.

Important Knowns and Recent Advances. Hormone receptors (eg, estrogen, testosterone, and relaxin) have been localized on the human ACL,^{186–190} suggesting they are capable of regulating gene expression and collagen metabolism in a way that may influence the biology of the ACL and other soft tissue structures. This idea is supported by studies that have demonstrated associations of normal physiologic variations in sex hormone concentrations across the menstrual cycle with substantial changes in markers of collagen metabolism and production,¹⁹¹ kneejoint laxity,^{130–135} muscle stiffness,¹³¹ and the muscle-stretch reflex.¹⁹² These biological changes may also have secondary neuromechanical consequences, as previously noted.^{155,156,158–160}

Although results from prior epidemiologic studies have suggested that the risk of an ACL injury appears to be greater during the preovulatory phase of the menstrual cycle compared with the postovulatory phase,^{193–197} our understanding of the underlying mechanism for this increased likelihood has not advanced. Understanding these processes is difficult given the substantial variability in the timing and magnitude of hormone changes among individuals,¹³⁴ the time-dependent effect for sex hormones and other remodeling agents to influence a change in ACL tissue characteristics,^{134,190} and the potential interactions among several sex hormones, secondary messengers, remodeling proteins, mechanical stresses, and genetic influences.* For example, interactions among hormones, mechanical stress, and altered ACL structure and metabolism have been observed in some animal models.^{148,205,206}

In recent years, the hormone relaxin has gained attention as a potential ACL injury risk factor based on a prospective study of National Collegiate Athletic Association Division I female athletes that showed elevated concentrations in those with ACL injuries compared with uninjured controls.²⁰⁷ Even though studies examining the effect of relaxin on collagen metabolism in human knee ligaments in situ are lacking, in vivo and in vitro animal studies and human cell culture studies suggest that relaxin administered at physiologic levels can have a profound effect on soft tissue remodeling.^{204,208–211} In turn, this may lead to a less organized²¹² and less dense (both in fiber diameter and density) collagen structure, 204,212 which may manifest as a weaker and more lax ACL.¹⁸⁶ This concept is supported by an animal model in which guinea pig ACLs treated with relaxin at pregnancy levels were 13% more lax and 36% to 49% weaker.²¹³

Unknowns and Directions for Future Research.

- 1. Our understanding of the biological consequences of sex hormones on collagen metabolism and the structural integrity of the ACL in the human knee in situ remains limited. The underlying sex-specific molecular and genetic mechanisms of sex hormones on ACL structure, metabolism, and mechanical properties and how mechanical stress on the ACL alters these relationships and injury risk potential remain important areas of study.
- 2. Good evidence demonstrates that some (but not all) females experience substantial cyclic changes in their laxity and knee-joint biomechanics across the menstrual cycle, ^{155,158,160} but it is not yet possible to clinically screen for these potentially high-risk individuals. Understanding the underlying processes that regulate ligament biology and resultant ligament behavior may allow us to better screen for these individuals and prospectively examine how these factors influence injury risk potential.
- 3. Given the time-dependent effect of sex hormones on soft tissue structures, we need to determine how the time of injury occurrence aligns with acute changes in ACL structure and metabolism or knee-laxity changes and how the rate of increase or the time durations of amplitude peaks in hormone fluctuations across the menstrual cycle play roles in the magnitude or timing of soft tissue changes.
- 4. When examining hormone influences in physically active females, it is important to match the complexity of interparticipant differences in timing, magnitude, and interactive changes in sex hormone concentrations across the cycle with our study designs. Investigations should (1) verify the phases of the cycle (or desired hormone environment) with actual hormone measurements (considering relevant hormones to include estrogen, progesterone, and possibly others) rather than relying on calendar day of the cycle²¹⁴ and (2) obtain multiple hormone samples over repeated days to better characterize hormone profiles within a given female.^{215,216} This may be particularly important when documenting relaxin exposure. Because relaxin rises and peaks during a relatively short window

*References 134, 173, 187, 190, 191, 198–204.

(6-10 days) after ovulation based on other hormonal events,^{217–220} we do not know if the high proportion of undetectable levels within a single sample (as high as $64\%-80\%)^{221-224}$ is due to timing or suggests that only a subset of females are exposed to appreciable relaxin levels due to mediating factors.^{219,221,225}

- 5. Because cyclic hormone concentrations affect soft tissues and knee-joint function, studies comparing females with males should be made during the early follicular phase in females, when hormone levels are at their nadirs (preferably 3–7 days postmenses).
- 6. Menstrual dysfunction in exercising females may be more prevalent than originally thought (up to 50% in exercising females).^{216,226,227} Thus, the effects of menstrual disturbances on injury risk potential are poorly understood.
- 7. Although the authors of epidemiologic studies have reported no protective effect of oral contraceptives on ACL injury risk,^{72,73} these medications are known to vary substantially in the potency and androgenicity of the progestin compound delivered, which ultimately determines the extent to which they counteract the estrogenic effects.²²⁸ We need to better understand how the different progestins influence soft tissue structures, knee function, and ACL injury risk. Furthermore, the exogenous hormones delivered in different types of contraceptives (eg, vaginal ring, transdermal patch, oral pills) are metabolized differently and may have different effects on musculoskeletal tissues.²²⁹ Relevant comparisons should then be made among users of hormonal contraceptives (of all 3 types) and eumenorrheic, amenorrheic, and oligomenorrheic females to determine if ACL injury risk or observed soft tissue changes vary.

SCREENING

For effective injury prevention, prospective risk factors for ACL injury should be established before preventive measures are introduced.²³⁰ Abnormal movement strategies may be modifiable and can be targeted in injury prevention.

Important Knowns and Recent Advances

Advances in clinic- or field-based screening tools may play an important role in identifying athletes who will benefit from injury-prevention programs.²⁶ Clinically oriented screening tools (eg, LESS, tuck jump) show good agreement with laboratory-based biomechanics (concurrent validity).^{231–234} Screening tools are sensitive in detecting changes in movement quality over time.^{235,236} Additionally, the ability of clinically oriented screening tools to identify individuals at risk for future ACL injury may be population specific (eg, sex, age, sport).^{26,27,233} For example, the LESS effectively identified elite youth soccer athletes at higher risk of sustaining ACL injuries, with a score of 5 yielding a sensitivity of 86% and a specificity of 64%.²⁶ However, the LESS has not been found to predict future injury in adults.²³⁷

Data continue to show that athletes are at higher risk of ACL injury in game situations: increased game-play exposure increases ACL injury risk.^{237,238} This suggests that screening in more of a real-life scenario may be of benefit in identifying those at greatest risk of injury.

Younger athletes with a previous ACL injury should be considered at higher risk for subsequent injury^{239–244} and identified through preseason screening, especially in pivoting and cutting sports.²⁴⁵ In his keynote summary, Paterno³ briefly reviewed the risk of subsequent injury.

Unknowns and Directions for Future Research

- 1. Which elements (eg, specific faulty movements, combination of faulty movements, strength assessment) of clinically oriented screening tools predict future ACL injury risk (predictive validity) is unknown. We need to develop other clinically oriented screening tools that have good sensitivity and specificity for predicting future ACL injury risk. Wearable sensors and advanced in-game monitoring may be helpful in assessing ACL injury risk and should be further explored.
- We must understand how laboratory-based (eg, drop vertical jump) and clinically oriented screening tools (eg, tuck jump, 2-dimensional assessment, clinical nomogram) predict ACL and other lower extremity injuries.^{25,26,246,247}
- 3. Although ongoing improvements to inexpensive and accurate screening methods to identify high-risk athletes remain of interest, enhancing existing training protocols and implementing neuromuscular training for all young athletes are warranted.²⁴⁸ Future authors should identify the relative effectiveness of injury-prevention programs that are seamlessly integrated into universal training and, potentially, physical education programs.
- 4. The most common tests used by professional male soccer clubs to identify noncontact ACL injury risk are the functional movement screen, questionnaires to recognize risk factors such as previous injury, and isokinetic assessments.²⁴⁹ Because the predictive ability of these tests is uncertain, clinicians must become familiar with evidence-based ACL injury-screening tools.
- 5. It is important to consider that when screening for characteristics that predict ACL injuries, identified and validated ACL injury risk factors may not align with more traditional variables that have been shown to directly increase ACL loading in cadaver-based or simulation studies (ie, anterior tibial shear, knee valgus, and internal rotation torque). Continued investigations to recognize ACL injury risk factors through clinical screens are necessary.⁵⁰

INJURY PREVENTION

Injury-prevention programs have been shown to reduce the incidence of ACL injuries. However, overall ACL injury rates and the associated sex disparity have not yet diminished. We still have much to learn in order to maximize the effectiveness of these programs, determine areas for improvement, and establish highly sensitive screening tools to target prevention efforts at those with the greatest risk for injury.

Important Knowns and Recent Advances

Various injury-prevention programs that incorporate elements of balance training, plyometric training, education, strengthening, and technique training or feedback have been shown to reduce ACL injury in females^{195,250–257} or alter biomechanical and neuromuscular variables thought to contribute to ACL injury.^{258–267} Injury-prevention programs with successful outcomes (eg, injury-rate reduction,

improvement in neuromuscular control or performance) are performed 2 to 3 times per week and last for a minimum of 10 to 15 minutes.[†]

Most exercises implemented in male soccer players to prevent noncontact injuries involve eccentric exercise (in general), balance, and specific hamstrings eccentric training.²⁴⁹ However, data regarding the effectiveness of prevention programs to reduce ACL injuries or modify risk factors for ACL injuries in male athletes are scarce and inconclusive as to recommendations for single-legged and multidirectional maneuvers.^{269,270}

Improvements in movement quality after 12 weeks of training do not appear to be retained once the program ends. Thus, optimal results for retention and transfer require training principles that entail sufficient practice, variation, and frequency.²³⁶ Recent researchers have focused on improving the delivery of injury-prevention programs. For example, feedback should emphasize successful performance and ignore less successful attempts; this benefits learning because of its positive motivational effects.²⁷¹ Programs that focus on correcting individualized techniques also appear to be successful.^{272–274} Real-time feedback has been used to positively affect landing biomechanics.275-278 However, expert modeling combined with self-modeling seems to be the most effective way to change landing biomechanics.^{263,274,279-281} Additionally, motor skills can be learned with an internal focus of attention (ie, on the movements themselves, as in "flex your knees when landing") or with an external focus of attention (ie, focus on the movement effect, as in "touch the target as you land").²⁸² Age-appropriate injury-prevention training programs can effectively modify biomechanics in children.^{235,283}

Compliance has a strong influence on the success of ACL injury-prevention programs.^{284,285} However, knowledge and beliefs may not affect program adherence,284,286 so other motivational factors must be considered. Coaches may be hesitant to implement an ACL injury-prevention program because they may feel it is "too much," not their primary interest,^{286–288} does not offer a relative advantage over their existing practices, does not align with their needs, or is too complex to implement in their setting.²⁸⁹ Yet having support from coaches and senior administrators is extremely important.²⁹⁰ Athletes are willing to perform a lower extremity injury-prevention program if data indicate the program could improve performance and lead to fewer injuries.²⁹¹ Athletic performance is a crucial part of injury prevention; coaches and others are more likely to adopt a program that does not conflict with performance.²⁹² Also, injury-prevention training can improve performance in vertical-jump height, strength, and running speed.^{268,273,274,293,294} Established frameworks could greatly help in organizing future research endeavors for largescale injury-prevention implementation efforts.^{295,296} On average, the implementation of a universal training program would save \$100 per player per season and would reduce the incidence of ACL injury from 3% to 1.1% per season.²⁴⁸

Unknowns and Directions for Future Research

1. Although ACL injury-prevention programs are successful in reducing the risk of ACL injury, the ideal combination of and emphasis on training components within these

†References 195, 250, 252–254, 258, 260–264, 268.

programs should be further identified.^{285,297–299} Research is necessary to determine which specific program elements are effective and necessary to reduce injury risk or promote biomechanical changes. For example, the preventive role of strengthening should be further examined.^{53,300–309}

- 2. We must ascertain if streamlining injury-prevention programs, thereby making them more palatable to the public, will improve compliance. We need to establish if virtual-reality, real-time–feedback, and portable technologies will improve the implementation of and compliance with screening and injury-prevention programs.²⁶ With new products that use video feedback, exergaming, and "gamified" scoring, commonly encountered barriers that prevent the training staff from implementing and adhering to ACL injury-prevention programs might be overcome. This innovative approach might also help us to reach the goal of making injury prevention in the future just another aspect of the sport.³¹⁰
- 3. We should continue to investigate how a participant's sex, age, skill level, and type of sport factor into the type and variety of exercises prescribed and technique training and feedback provided. Specifically, the most important age or stage of growth and development to begin implementing injury-prevention training should be determined.
- 4. During injury-prevention training programs, technique training and feedback are frequently provided to improve movement patterns. An external focus seems to be advantageous over an internal focus; however, more research is needed to establish the most effective training protocols (eg, frequency, timing, combinations of focus of attention) for improving movement patterns and optimizing the transfer of these learned patterns to sport-specific movements performed during a game. For instance, consciously learning optimal movement patterns rehearsed during training sessions might result in suboptimal transfer to the field, where complex, unanticipated automatic movements are required.²⁸² Learning movement patterns with an external focus might enhance automatic motor control.³¹¹
- 5. Rehabilitation after ACL reconstruction that combines internal and external focus may enhance automatic movement control and improve performance.^{312–314} However, we need to investigate the effects of injury-prevention training on ACL injury rates in those with a history of ACL injury.
- 6. We must understand how injury-prevention programs influence lower extremity injuries other than ACL injury.
- 7. We need to continue to study the performance-enhancement benefits associated with regular injury-prevention training.
- 8. The modes and frequency of feedback and instruction should be further investigated. Self-controlled feedback enhances retention of movement patterns over time.²⁷⁴ Additionally, giving the athlete some control over a practice session (ie, an active role in deciding when to receive feedback) may enhance motor-skill learning when compared with prescribed training schedules.^{315,316} It is interesting that athletes often have a relatively strong sense of how well they perform.³¹⁷ This so-called self-controlled practice has generally been assumed to initiate a more active involvement of the athlete, enhancing

motivation and increasing the effort invested in practice.²⁸² In addition, sex-specific feedback strategies should be considered when implementing ACL injuryprevention programs. Males respond well to visual feedback, whereas females may prefer different learning strategies (eg, combined visual and verbal feedback).^{273,274}

- 9. Although well-controlled ACL injury-prevention programs reduce the incidence of ACL injuries, we have yet to effectively implement multifaceted programs in different settings that are sustainable over time (widespread implementation with high compliance and retention rates over the long term). Developing packaged preventive training programs that can be implemented broadly across different settings through appropriately educated and trained coaches or team leaders may facilitate compliance and efficacy. To that end, the following should be considered when developing largescale injury-prevention programs in the future: (1) lowcost, brief-time, packaged interventions; (2) adaptation of the program based on contextual factors for that setting (eg. sport, age, sex, environment); (3) incorporation of lay people (eg, coaches instead of athletic trainers or strength and conditioning specialists) to implement the program for that setting and population; (4) educating and obtaining organizational "buy in" from all levels (eg, school, club, administrators, coaches, players, parents); (5) embedding programs within the existing system when possible (eg, part of the warm-up or conditioning program, team challenge); and (6) developing written policies and procedures (ie, specifics of program, when to perform, how often to perform).
- 10. Although investigators have elucidated the influence of anatomical and structural factors on weight-bearing knee-joint neuromechanics,^{116,119,120,153,154,160} this information has yet to be incorporated in our ACL injury-prevention efforts.

ACKNOWLEDGMENTS

We thank the North Carolina Biotechnology Center (Research Triangle Park); The MotionMonitor (Chicago, IL); Biodex Medical Systems, Inc (Shirley, NY); Qualisys Motion Capture Systems (Gothenburg, Sweden); and the University of North Carolina at Greensboro Program for the Advancement of Girls and Women in Sport and Physical Activity for their support of the meeting.

REFERENCES

- Beynnon BD, Sturnick DR, Argentieri EC, et al. A sex-stratified multivariate risk factor model for anterior cruciate ligament injury. *J Athl Train.* 2015;50(10):1094–1096.
- Swanik CB. Brains and sprains: the brain's role in noncontact anterior cruciate ligament injuries. J Athl Train. 2015;50(10):1100– 1102.
- Paterno MV. Incidence and predictors of second anterior cruciate ligament injury after primary reconstruction and return to sport. J Athl Train. 2015;50(10):1097–1099.
- Shultz SJ, Schmitz RJ, Benjaminse A, Chaudhari AM, Collins M, Padua DA. ACL Research Retreat VI: an update on ACL injury risk and prevention. *J Athl Train*. 2012;47(5):591–603.
- 5. Oh YK, Kreinbrink JL, Wojtys EM, Ashton-Miller JA. Effect of axial tibial torque direction on ACL relative strain and strain rate in

an in vitro simulated pivot landing. J Orthop Res. 2012;30(4):528–534.

- Oh YK, Lipps DB, Ashton-Miller JA, Wojtys EM. What strains the anterior cruciate ligament during a pivot landing? *Am J Sports Med.* 2012;40(3):574–583.
- Shin CS, Chaudhari AM, Andriacchi TP. The effect of isolated valgus moments on ACL strain during single-leg landing: a simulation study. *J Biomech.* 2009;42(3):280–285.
- Shin CS, Chaudhari AM, Andriacchi TP. Valgus plus internal rotation moments increase anterior cruciate ligament strain more than either alone. *Med Sci Sports Exerc.* 2011;43(8):1484–1491.
- Withrow TJ, Huston LJ, Wojtys EM, Ashton-Miller JA. The effect of an impulsive knee valgus moment on in vitro relative ACL strain during a simulated jump landing. *Clin Biomech (Bristol, Avon)*. 2006;21(9):977–983.
- Kiapour AM, Quatman CE, Goel VK, Wordeman SC, Hewett TE, Demetropoulos CK. Timing sequence of multi-planar knee kinematics revealed by physiologic cadaveric simulation of landing: implications for ACL injury mechanism. *Clin Biomech (Bristol, Avon)*. 2014;29(1):75–82.
- Markolf KL, Jackson SR, Foster B, McAllister DR. ACL forces and knee kinematics produced by axial tibial compression during a passive flexion-extension cycle. J Orthop Res. 2014;32(1):89–95.
- Arms SW, Pope MH, Johnson RJ, Fischer RA, Arvidsson I, Eriksson E. The biomechanics of anterior cruciate ligament rehabilitation and reconstruction. *Am J Sports Med.* 1984;12(1):8– 18.
- Berns GS, Hull ML, Patterson HA. Strain in the anteromedial bundle of the anterior cruciate ligament under combination loading. *J Orthop Res.* 1992;10(2):167–176.
- Levine JW, Kiapour AM, Quatman CE, et al. Clinically relevant injury patterns after an anterior cruciate ligament injury provide insight into injury mechanisms. *Am J Sports Med.* 2013;41(2):385– 395.
- Chandrashekar N, Bakker R, Tomescu S, Brennan E, Hangalur G, Laing A. The effect of sagittal-plane mechanics on ACL strain during jump landing [abstract]. J Athl Train. 50(10):1107.
- Shimokochi Y, Shultz SJ. Mechanisms of noncontact anterior cruciate ligament injury. J Athl Train. 2008;43(4):396–408.
- Koga H, Bahr R, Myklebust G, Engebretsen L, Grund T, Krosshaug T. Estimating anterior tibial translation from model-based imagematching of a noncontact anterior cruciate ligament injury in professional football: a case report. *Clin J Sport Med.* 2011;21(3): 271–274.
- Koga H, Nakamae A, Shima Y, et al. Mechanisms for noncontact anterior cruciate ligament injuries: knee joint kinematics in 10 injury situations from female team handball and basketball. *Am J Sports Med.* 2010;38(11):2218–2225.
- Boden BP, Dean GS, Feagin JA Jr, Garrett WE Jr. Mechanisms of anterior cruciate ligament injury. *Orthopedics*. 2000;23(6):573–578.
- Waldén M, Krosshaug T, Bjørneboe J, Andersen TE, Faul O, Hägglund M. Three distinct mechanisms predominate in noncontact anterior cruciate ligament injuries in male professional football players: a systematic video analysis of 39 cases. *Br J Sports Med.* In press.
- Krosshaug T, Nakamae A, Boden BP, et al. Mechanisms of anterior cruciate ligament injury in basketball: video analysis of 39 cases. *Am J Sports Med.* 2007;35(3):359–367.
- 22. Olsen OE, Myklebust G, Engebretsen L, Bahr R. Injury mechanisms for anterior cruciate ligament injuries in team handball: a systematic video analysis. *Am J Sports Med.* 2004;32(4):1002–1012.
- 23. Hewett TE, Torg JS, Boden BP. Video analysis of trunk and knee motion during non-contact anterior cruciate ligament injury in

female athletes: lateral trunk and knee abduction motion are combined components of the injury mechanism. *Br J Sports Med.* 2009;43(6):417–422.

- Sheehan FT, Sipprell WH III, Boden BP. Dynamic sagittal plane trunk control during anterior cruciate ligament injury. *Am J Sports Med.* 2012;40(5):1068–1074.
- 25. Hewett TE, Myer GD, Ford KR, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med.* 2005;33(4):492–501.
- Padua DA, DiStefano LJ, Beutler AI, de la Motte SJ, DiStefano MJ, Marshall SW. The Landing Error Scoring System as a screening tool for an anterior cruciate ligament injury-prevention program in elite-youth soccer athletes. *J Athl Train*. 2015;50(6):589–595.
- Smith HC, Johnson RJ, Shultz SJ, et al. A prospective evaluation of the Landing Error Scoring System (LESS) as a screening tool for anterior cruciate ligament injury risk. *Am J Sports Med.* 2012;40(3): 521–526.
- Blackburn JT, Padua DA. Sagittal-plane trunk position, landing forces, and quadriceps electromyographic activity. J Athl Train. 2009;44(2):174–179.
- 29. Shimokochi Y, Yong Lee S, Shultz SJ, Schmitz RJ. The relationships among sagittal-plane lower extremity moments: implications for landing strategy in anterior cruciate ligament injury prevention. *J Athl Train.* 2009;44(1):33–38.
- Cerulli G, Benoit DL, Lamontagne M, Caraffa A, Liti A. In vivo anterior cruciate ligament strain behaviour during a rapid deceleration movement: case report. *Knee Surg Sports Traumatol Arthrosc.* 2003;11(5):307–311.
- Cassidy K, Hangalur G, Sabharwal P, Chandrashekar N. Combined in vivo/in vitro method to study anteriomedial bundle strain in the anterior cruciate ligament using a dynamic knee simulator. J Biomech Eng. 2013;135(3):35001.
- Myers CA, Torry MR, Shelburne KB, et al. In vivo tibiofemoral kinematics during 4 functional tasks of increasing demand using biplane fluoroscopy. *Am J Sports Med.* 2012;40(1):170–178.
- Schmitz RJ, Kim H, Shultz SJ. Effect of axial load on anterior tibial translation when transitioning from non-weight bearing to weight bearing. *Clin Biomech (Bristol, Avon)*. 2010;25(1):77–82.
- Decker MJ, Torry MR, Wyland DJ, Sterett WI, Richard SJ. Gender differences in lower extremity kinematics, kinetics and energy absorption during landing. *Clin Biomech (Bristol, Avon)*. 2003; 18(7):662–669.
- Houck JR, Duncan A, De Haven KE. Comparison of frontal plane trunk kinematics and hip and knee moments during anticipated and unanticipated walking and side step cutting tasks. *Gait Posture*. 2006;24(3):314–322.
- Pollard CD, Sigward SM, Powers CM. Gender differences in hip joint kinematics and kinetics during side-step cutting maneuver. *Clin J Sport Med.* 2007;17(1):38–42.
- Schmitz RJ, Kulas AS, Perrin DH, Riemann BL, Shultz SJ. Sex differences in lower extremity biomechanics during single leg landings. *Clin Biomech (Bristol, Avon)*. 2007;22(6):681–688.
- Utturkar GM, Irribarra LA, Taylor KA, et al. The effects of a valgus collapse knee position on in vivo ACL elongation. *Ann Biomed Eng.* 2013;41(1):123–130.
- Chappell JD, Herman DC, Knight BS, Kirkendall DT, Garrett WE, Yu B. Effect of fatigue on knee kinetics and kinematics in stopjump tasks. *Am J Sports Med.* 2005;33(7):1022–1029.
- Cortes N, Quammen D, Lucci S, Greska E, Onate J. A functional agility short-term fatigue protocol changes lower extremity mechanics. *J Sports Sci.* 2012;30(8):797–805.

- Kernozek TW, Torry MR, Iwasaki M. Gender differences in lower extremity landing mechanics caused by neuromuscular fatigue. *Am J Sports Med.* 2008;36(3):554–565.
- McLean SG, Fellin RE, Suedekum N, Calabrese G, Passerallo A, Joy S. Impact of fatigue on gender-based high-risk landing strategies. *Med Sci Sports Exerc*. 2007;39(3):502–514.
- Quammen D, Cortes N, Van Lunen BL, Lucci S, Ringleb SI, Onate J. Two different fatigue protocols and lower extremity motion patterns during a stop-jump task. *J Athl Train*. 2012; 47(1):32–41.
- Schmitz RJ, Cone JC, Tritsch AJ, et al. Changes in drop-jump landing biomechanics during prolonged intermittent exercise. *Sports Health.* 2014;6(2):128–135.
- 45. Raja Azidin RM, Sankey S, Drust B, Robinson MA, Vanrenterghem J. Effects of treadmill versus overground soccer match simulations on biomechanical markers of anterior cruciate ligament injury risk in side cutting. *J Sports Sci.* 2015;33(13):1332–1341.
- Agel J, Arendt EA, Bershadsky B. Anterior cruciate ligament injury in National Collegiate Athletic Association basketball and soccer: a 13-year review. *Am J Sports Med.* 2005;33(4):524–530.
- Joseph AM, Collins CL, Henke NM, Yard EE, Fields SK, Comstock RD. A multisport epidemiologic comparison of anterior cruciate ligament injuries in high school athletics. *J Athl Train*. 2013;48(6): 810–817.
- Carson DW, Ford KR. Sex differences in knee abduction during landing: a systematic review. *Sports Health*. 2011;3(4):373–382.
- Beaulieu ML, McLean SG. Sex-dimorphic landing mechanics and their role within the noncontact ACL injury mechanism: evidence, limitations and directions. *Sports Med Arthrosc Rehabil Ther Technol.* 2012;4(10):1–13.
- Zazulak BT, Hewett TE, Reeves NP, Goldberg B, Cholewicki J. Deficits in neuromuscular control of the trunk predict knee injury risk: a prospective biomechanical-epidemiologic study. *Am J Sports Med.* 2007;35(7):1123–1130.
- Kulas AS, Zalewski P, Hortobagyi T, DeVita P. Effects of added trunk load and corresponding trunk position adaptations on lower extremity biomechanics during drop-landings. *J Biomech.* 2008; 41(1):180–185.
- McLean SG, Huang X, Van Den Bogert AJ. Association between lower extremity posture at contact and peak knee valgus moment during sidestepping: implications for ACL injury. *Clin Biomech* (*Bristol, Avon*). 2005;20(8):863–870.
- Frank B, Bell DR, Norcross MF, Blackburn JT, Goerger BM, Padua DA. Trunk and hip biomechanics influence anterior cruciate loading mechanisms in physically active participants. *Am J Sports Med.* 2013;41(11):2676–2683.
- Sigward SM, Powers CM. Loading characteristics of females exhibiting excessive valgus moments during cutting. *Clin Biomech* (*Bristol, Avon*). 2007;22(7):827–833.
- Jamison ST, Pan X, Chaudhari AM. Knee moments during run-tocut maneuvers are associated with lateral trunk positioning. J Biomech. 2012;45(11):1881–1885.
- Swanik CB, Covassin T, Stearne DJ, Schatz P. The relationship between neurocognitive function and noncontact anterior cruciate ligament injuries. *Am J Sports Med.* 2007;35(6):943–948.
- Herman DC, Barth JT. The influence of neurocognitive performance on trunk stabilities varies with sex [abstract]. *J Athl Train*. 2015;50(10):1105–1106.
- McLean SG, Mallett KF, Arruda EM. Deconstructing the anterior cruciate ligament: what we know and do not know about function, material properties, and injury mechanics. *J Biomech Eng.* 2015; 137(2):020906.

- Quatman CE, Quatman CC, Hewett TE. Prediction and prevention of musculoskeletal injury: a paradigm shift in methodology. *Br J Sports Med.* 2009;43(14):1100–1107.
- Kristianslund E, Krosshaug T. Comparison of drop jumps and sportspecific sidestep cutting: implications for anterior cruciate ligament injury risk screening. *Am J Sports Med.* 2013;41(3):684–688.
- Ford KR, Taylor JB, Nguyen A. Kinematic landing differences following biofeedback training in female athletes [abstract]. J Athl Train. 2015;50(10):1111.
- Kulas AS, Hortobagyi T, DeVita P. Trunk position modulates anterior cruciate ligament forces and strains during a single-leg squat. *Clin Biomech (Bristol, Avon)*. 2012;27(1):16–21.
- 63. Escamilla RF, Macleod TD, Wilk KE, Paulos L, Andrews JR. Anterior cruciate ligament strain and tensile forces for weightbearing and non-weight-bearing exercises: a guide to exercise selection. J Orthop Sports Phys Ther. 2012;42(3):208–220.
- Decker LM, Moraiti C, Stergiou N, Georgoulis AD. New insights into anterior cruciate ligament deficiency and reconstruction through the assessment of knee kinematic variability in terms of nonlinear dynamics. *Knee Surg Sports Traumatol Arthrosc.* 2011; 19(10):1620–1633.
- Pataky TC, Robinson MA, Vanrenterghem J. Vector field statistical analysis of kinematic and force trajectories. *J Biomech.* 2013; 46(14):2394–2401.
- Pataky TC, Vanrenterghem J, Robinson MA. Two-way ANOVA for scalar trajectories, with experimental evidence of non-phasic interactions. J Biomech. 2015;48(1):186–189.
- Robinson MA, Donnelly CJ, Vanrenterghem J, Pataky TC. Vectorfield statistical analysis reveals important insight into knee joint moments during sidestepping [abstract]. *J Athl Train*. 2015;50(10): 1107–1108.
- Baumeister J, von Detten S, van Niekerk SM, Schubert M, Ageberg E, Louw QA. Brain activity in predictive sensorimotor control for landings: an EEG pilot study. *Int J Sports Med.* 2013;34(12):1106–1111.
- Lipps DB, Wojtys EM, Ashton-Miller JA. Anterior cruciate ligament fatigue failures in knees subjected to repeated simulated pivot landings. *Am J Sports Med.* 2013;41(5):1058–1066.
- Stearns KM, Powers CM. Improvements in hip muscle performance result in increased use of the hip extensors and abductors during a landing task. *Am J Sports Med.* 2014;42(3):602–609.
- Stearns KM, Keim RG, Powers CM. Influence of relative hip and knee extensor muscle strength on landing biomechanics. *Med Sci Sports Exerc*. 2013;45(5):935–941.
- Montgomery MM, Shultz SJ, Schmitz RJ, Wideman L, Henson RA. Influence of lean body mass and strength on landing energetics. *Med Sci Sports Exerc*. 2012;44(12):2376–2383.
- Schmitz RJ, Shultz SJ. Contribution of knee flexor and extensor strength on sex-specific energy absorption and torsional joint stiffness during drop jumping. J Athl Train. 2010;45(5):445–452.
- Shultz SJ, Nguyen AD, Leonard MD, Schmitz RJ. Thigh strength and activation as predictors of knee biomechanics during a drop jump task. *Med Sci Sports Exerc.* 2009;41(4):857–866.
- Weinhandl JT, Earl-Boehm JE, Ebersole KT, Huddleston WE, Armstrong BS, O'Connor KM. Reduced hamstring strength increases anterior cruciate ligament loading during anticipated sidestep cutting. *Clin Biomech (Bristol, Avon)*. 2014;29(7):752–759.
- Barber-Westin SD, Galloway M, Noyes FR, Corbett G, Walsh C. Assessment of lower limb neuromuscular control in prepubescent athletes. *Am J Sports Med.* 2005;33(12):1853–1860.
- Barber-Westin SD, Noyes FR, Galloway M. Jump-land characteristics and muscle strength development in young athletes: a gender

comparison of 1140 athletes 9 to 17 years of age. *Am J Sports Med.* 2006;34(3):375–384.

- Hass CJ, Schick EA, Tillman MD, Chow JW, Brunt D, Cauraugh JH. Knee biomechanics during landings: comparison of pre- and postpubescent females. *Med Sci Sports Exerc.* 2005;37(1):100–107.
- Hewett TE, Myer GD, Ford KR. Decrease in neuromuscular control about the knee with maturation in female athletes. *J Bone Joint Surg Am.* 2004;86:1601–1608.
- Quatman CE, Ford KR, Myer GD, Hewett TE. Maturation leads to gender differences in landing force and vertical jump performance: a longitudinal study. *Am J Sports Med.* 2006;34(5):806–813.
- Russell PJ, Croce RV, Swartz EE, Decoster LC. Knee-muscle activation during landings: developmental and gender comparisons. *Med Sci Sports Exerc.* 2007;39(1):159–170.
- Swartz EE, Decoster LC, Russell PJ, Croce RV. Effects of developmental stage and sex on lower extremity kinematics and vertical ground reaction forces during landing. *J Athl Train*. 2005; 40(1):9–14.
- Ford KR, Shapiro R, Myer GD, VanDenBogert AJ, Hewett TE. Longitudinal sex differences during landing in knee abduction in young athletes. *Med Sci Sports Exerc.* 2010;42(10):1923–1931.
- Hewett TE, Lynch TR, Myer GD, Ford KR, Gwin RC, Heidt RS Jr. Multiple risk factors related to familial predisposition to anterior cruciate ligament injury: fraternal twin sisters with anterior cruciate ligament ruptures. *Br J Sports Med.* 2010;44(12):848–855.
- Sigward SM, Pollard CD, Havens KL, Powers CM. The influence of sex and maturation on knee mechanics during side-step cutting. *Med Sci Sports Exerc.* 2012;44(8):1497–1503.
- Kim KW, Lim BO. Effects of menarcheal age on the anterior cruciate ligament injury risk factors during single-legged drop landing in female artistic elite gymnasts. *Arch Orthop Trauma Surg.* 2014;134(11):1565–1571.
- Hewett TE, Myer GD, Kiefer AW, Ford KR. Longitudinal increases in knee abduction moments in females during adolescent growth [published online ahead of print May 12, 2015]. *Med Sci Sports Exerc.* doi:10.1249/MSS.000000000000700.
- Holden S, Boreham C, Delahunt E. Sex differences in landing biomechanics and postural stability of adolescent athletes: a systematic review [abstract]. *J Athl Train*. 2015;50(10):1106.
- Malfait B, Sankey S, Firhad Raja Azidin RM, et al. How reliable are lower-limb kinematics and kinetics during a drop vertical jump? *Med Sci Sports Exerc.* 2014;46(4):678–685.
- Sankey SP, Raja Azidin RM, Robinson MA, et al. How reliable are knee kinematics and kinetics during side-cutting manoeuvres? *Gait Posture*. 2015;14(4):905–911.
- Myer GD, Bates NA, DiCesare CA, et al. Reliability of 3dimensional measures of single leg landing across three different institutions: implications for multi-center biomechanical and epidemiological research for secondary ACL injury prevention. J Sport Rehabil. 2015;24(2):198–209.
- Kristianslund E, Krosshaug T, van den Bogert AJ. Effect of low pass filtering on joint moments from inverse dynamics: implications for injury prevention. *J Biomech.* 2012;45(4):666–671.
- Roewer BD, Ford KR, Myer GD, Hewett TE. The "impact" of force filtering cut-off frequency on the peak knee abduction moment during landing: artefact or "artifiction"? *Br J Sports Med.* 2014; 48(6):464–468.
- Chaudhari AM, Zelman EA, Flanigan DC, Kaeding CC, Nagaraja HN. Anterior cruciate ligament-injured subjects have smaller anterior cruciate ligaments than matched controls: a magnetic resonance imaging study. *Am J Sports Med.* 2009;37(7):1282–1287.
- 95. Whitney DC, Sturnick DR, Vacek PM, et al. Relationship between the risk of suffering a first-time noncontact ACL injury and

geometry of the femoral notch and ACL: a prospective cohort study with a nested case-control analysis. *Am J Sports Med.* 2014;42(8): 1796–1805.

- Khan MS, Seon JK, Song EK. Risk factors for anterior cruciate ligament injury: assessment of tibial plateau anatomic variables on conventional MRI using a new combined method. *Int Orthop.* 2011; 35(8):1251–1256.
- Hashemi J, Chandrashekar N, Mansouri H, et al. Shallow medial tibial plateau and steep medial and lateral tibial slopes: new risk factors for anterior cruciate ligament injuries. *Am J Sports Med.* 2010;38(1):54–62.
- Everhart JS, Flanigan DC, Simon RA, Chaudhari AM. Association of non-contact ACL injury with presence and thickness of a bony ridge on the anteromedial aspect of the femoral intercondylar notch. *Am J Sports Med.* 2010;38(8):1667–1676.
- Stijak L, Herzog RF, Schai P. Is there an influence of the tibial slope of the lateral condyle on the ACL lesion? A case-control study. *Knee Surg Sports Traumatol Arthrosc.* 2008;16(2):112–117.
- 100. Beynnon BD, Hall JS, Sturnick DR, et al. Increased slope of the lateral tibial plateau subchondral bone is associated with greater risk of noncontact ACL injury in females but not in males: a prospective cohort study with a nested, matched case-control analysis. *Am J Sports Med.* 2014;42(5):1039–1048.
- LaPrade RF, Burnett QM. Femoral intercondylar notch stenosis and correlation to anterior cruciate ligament injuries. *Am J Sports Med.* 1994;22(2):198–302.
- 102. Shelbourne KD, Davis TJ, Klootwyk TE. The relationship between intercondylar notch width of the femur and the incidence of anterior cruciate ligament tears: a prospective study. *Am J Sports Med.* 1998; 26(3):402–408.
- Souryal TO, Freeman TR. Intercondylar notch size and anterior cruciate ligament injuries in athletes. *Am J Sports Med.* 1993;21(4): 535–539.
- 104. Uhorchak JM, Scoville CR, Williams GN, Arciero RA, St Pierre P, Taylor DC. Risk factors associated with non-contact injury of the anterior cruciate ligament. *Am J Sports Med.* 2003;31(6):831– 842.
- 105. Ireland ML, Ballantyne BT, Little K, McClay IS. A radiographic analysis of the relationship between the size and shape of the intercondylar notch and anterior cruciate ligament injury. *Knee Surg Sports Traumatol Arthrosc.* 2001;9(4):200–205.
- Souryal TO, Moore HA, Evans JP. Bilaterality in anterior cruciate ligament injuries. Am J Sports Med. 1988;16(5):449–454.
- 107. Domzalski M, Grzelak P, Gabos P. Risk factors for anterior cruciate ligament injury in skeletally immature patients: analysis of intercondylar notch width using magnetic resonance imaging. *Int Orthop.* 2010;34(5):703–707.
- 108. Zeng C, Gao S, Wei J, et al. The influence of the intercondylar notch dimensions on injury of the anterior cruciate ligament: a meta-analysis. *Knee Surg Sports Traumatol Arthrosc.* 2013;21(4): 804–815.
- 109. Simon RA, Everhart JS, Nagaraja HN, Chaudhari AM. A casecontrol study of anterior cruciate ligament volume, tibial plateau slopes and intercondylar notch dimensions in ACL-injured knees. J Biomech. 2010;43(9):1702–1707.
- 110. Sturnick DR, Vacek PM, DeSarno MJ, et al. Combined anatomic factors predicting risk of anterior cruciate ligament injury for males and females. *Am J Sports Med.* 2015;43(4):839–847.
- 111. Chandrashekar N, Slauterbeck J, Hashemi J. Sex-based differences in the anthropometric characteristics of the anterior cruciate ligament and its relation to intercondylar notch geometry. *Am J Sports Med.* 2005;33(10):1492–1498.

- 112. Hashemi J, Chandrashekar N, Mansouri H, Slauterbeck J, Hardy DM. The human anterior cruciate ligament: sex differences in ultrastructure and correlation with biomechanical properties. J Orthop Res. 2008;26(7):945–950.
- 113. Chandrashekar N, Mansour JM, Slauterbeck J, Hashemi J. Sexbased differences in the tensile properties of the human anterior cruciate ligament. *J Biomech*. 2006;39(16):2943–2950.
- 114. Hashemi J, Chandrashekar N, Gill B, et al. The geometry of the tibial plateau and its influence on the biomechanics of the tibiofemoral joint. *J Bone Joint Surg Am.* 2008;90(12):2724–2734.
- 115. Hudek R, Fuchs B, Regenfelder F, Koch PP. Is noncontact ACL injury associated with the posterior tibial and meniscal slope? *Clin Orthop Relat Res.* 2011;469(8):2377–2384.
- 116. McLean SG, Lucey SM, Rohrer S, Brandon C. Knee joint anatomy predicts high-risk in vivo dynamic landing knee biomechanics. *Clin Biomech (Bristol, Avon)*. 2010;25(8):781–788.
- Dejour H, Bonnin M. Tibial translation after anterior cruciate ligament rupture: two radiological tests compared. J Bone Joint Surg Br. 1994;76(5):745–749.
- Giffin JR, Vogrin TM, Zantop T, Woo SL, Harner CD. Effects of increasing tibial slope on the biomechanics of the knee. *Am J Sports Med.* 2004;32(2):376–382.
- 119. McLean SG, Oh YK, Palmer ML, et al. The relationship between anterior tibial acceleration, tibial slope, and ACL strain during a simulated jump landing task. *J Bone Joint Surg Am.* 2011;93(14): 1310–1317.
- 120. Lipps DB, Oh YK, Ashton-Miller JA, Wojtys EM. Morphologic characteristics help explain the gender difference in peak anterior cruciate ligament strain during a simulated pivot landing. *Am J Sports Med.* 2012;40(1):32–40.
- Shultz SJ, Schmitz RJ. Tibial plateau geometry influences lower extremity biomechanics during landing. *Am J Sports Med.* 2012; 40(9):2029–2036.
- 122. Scerpella TA, Stayer TJ, Makhuli BZ. Ligamentous laxity and noncontact anterior cruciate ligament tears: a gender based comparison. *Orthopaedics*. 2005;28(7):656–660.
- 123. Woodford-Rogers B, Cyphert L, Denegar CR. Risk factors for anterior cruciate ligament injury in high school and college athletes. *J Athl Train*. 1994;29(4):343–346.
- 124. Kramer LC, Denegar CR, Buckley WE, Hertel J. Factors associated with anterior cruciate ligament injury: history in female athletes. J Sports Med Phys Fitness. 2007;47(4):446–454.
- Loudon JK, Jenkins W, Loudon KL. The relationship between static posture and ACL injury in female athletes. J Orthop Sports Phys Ther. 1996;24(2):91–97.
- 126. Myer GD, Ford KR, Paterno MV, Nick TG, Hewett TE. The effects of generalized joint laxity on risk of anterior cruciate ligament injury in young female athletes. *Am J Sports Med.* 2008;36(6): 1073–1080.
- 127. Ramesh R, VonArx O, Azzopardi T, Schranz PJ. The risk of anterior cruciate ligament rupture with generalised joint laxity. J Bone Joint Surg Br. 2005;87(6):800–803.
- 128. Branch TP, Browne JE, Campbell JD, et al. Rotational laxity greater in patients with contralateral anterior cruciate ligament injury than healthy volunteers. *Knee Surg Sports Traumatol Arthrosc.* 2010; 18(10):1379–1384.
- 129. Bell RD, Shultz SJ, Wideman L, Henrich VC. Collagen gene variants previously associated with ACL injury risk are also associated with joint laxity. *Sports Health*. 2012;4(4):312–318.
- Deie M, Sakamaki Y, Sumen Y, Urabe Y, Ikuta Y. Anterior knee laxity in young women varies with their menstrual cycle. *Int Orthop.* 2002;26(3):154–156.

- 131. Eiling W, Bryant AL, Petersen W, Murphy A, Hohmann E. Effects of menstrual cycle hormone fluctuations on musculoskeletal stiffness and knee joint laxity. *Knee Surg Sports Traumatol Arthrosc.* 2007;15(2):126–132.
- Heitz NA. Hormonal changes throughout the menstrual cycle and increased anterior cruciate ligament laxity in females. *J Athl Train*. 1999;343(2):144–149.
- 133. Shultz SJ, Levine BJ, Nguyen AD, Kim HS, Montgomery MM, Perrin DH. A comparison of cyclic variations in anterior knee laxity, genu recurvatum and general joint laxity across the menstrual cycle. *J Orthop Res.* 2010;28(11):1411–1417.
- Shultz SJ, Sander TC, Kirk SE, Johnson M, Perrin DH. Relationship between sex hormones and anterior knee laxity across the menstrual cycle. *Med Sci Sports Exerc*. 2004;36(7):1165–1174.
- 135. Shultz SJ, Schmitz RJ, Beynnon BD. Variations in varus/valgus and internal/external rotational knee laxity and stiffness across the menstrual cycle. J Orthop Res. 2011;29(3):318–325.
- Shultz SJ, Dudley WN, Kong Y. Identifying knee laxity profiles and associated physical characteristics. J Athl Train. 2012;47(2):159– 169.
- 137. Shultz SJ, Pye ML, Montgomery MM, Schmitz RJ. Associations between lower extremity muscle mass and multiplanar knee laxity and stiffness: a potential explanation for sex differences in frontal and transverse plane knee laxity. *Am J Sports Med.* 2012;40(12): 2836–2844.
- Wang HM, Kulas AS, Kraft RA, Shultz SJ, Schmitz RJ. Quadriceps muscle volume is predictive of ACL volume [abstract]. *J Athl Train*. 2015;50(10):1105.
- 139. Rozzi SL, Lephart SM, Gear WS, Fu FH. Knee joint laxity and neuromuscular characteristics of male and female soccer and basketball players. *Am J Sports Med.* 1999;27(3):312–319.
- Shultz SJ, Kirk SE, Sander TC, Perrin DH. Sex differences in knee laxity change across the female menstrual cycle. *J Sports Med Phys Fitness*. 2005;45(4):594–603.
- Beynnon BD, Bernstein I, Belisle A, et al. The effect of estradiol and progesterone on knee and ankle joint laxity. *Am J Sports Med.* 2005;33(9):1298–1304.
- Nguyen AD, Shultz SJ. Sex differences in lower extremity posture. J Orthop Sports Phys Ther. 2007;37(7):389–398.
- 143. Trimble MH, Bishop MD, Buckley BD, Fields LC, Rozea GD. The relationship between clinical measurements of lower extremity posture and tibial translation. *Clin Biomech (Bristol, Avon)*. 2002; 17(4):286–290.
- 144. Hsu W, Fisk JA, Yamamoto Y, Debski RE, Woo SLY. Differences in torsional joint stiffness of the knee between genders: a human cadaveric study. *Am J Sports Med.* 2006;34(5):765–770.
- 145. Markolf KL, Graff-Radford A, Amstutz HC. In vivo knee stability: a quantitative assessment using an instrumented clinical testing apparatus. *J Bone Joint Surg Am.* 1978;60(5):664–674.
- 146. Sharma L, Lou C, Felson DT, et al. Laxity in healthy and osteoarthritic knees. *Arthritis Rheum.* 1999;42(5):861–870.
- 147. Shultz SJ, Schmitz RJ, Cone JR, et al. Multiplanar knee laxity increases during a 90-min intermittent exercise protocol. *Med Sci Sports Exerc*. 2013;45(8):1553–1561.
- 148. Comerford EJ, Tarlton JF, Innes JF, Johnson KA, Amis AA, Bailey AJ. Metabolism and composition of the canine anterior cruciate ligament relate to differences in knee joint mechanics and predisposition to ligament rupture. *J Orthop Res.* 2005;23(1): 61–66.
- 149. Quasnichka HL, Anderson-MacKenzie JM, Tarlton JF, Sims TJ, Billingham MEJ, Bailey AJ. Cruciate ligament laxity and femoral intercondylar notch narrowing in early-stage knee osteoarthritis. *Arthritis Rheum*. 2005;52(10):3100–3109.

- 150. Wang VM, Banack TM, Tsai CW, Flatow EL, Jepsen KJ. Variability in tendon and knee joint biomechanics among inbred mouse strains. *J Orthop Res.* 2006;24(6):1200–1207.
- Shultz SJ, Carcia CR, Perrin DH. Knee joint laxity affects muscle activation patterns in the healthy knee. *J Electromyogr Kinesiol*. 2004;14(4):475–483.
- 152. Shultz SJ, Shimokochi Y, Nguyen A, et al. Non-weight bearing anterior knee laxity is related to anterior tibial translation during transition from non-weight bearing to weight bearing. *J Orthop Res.* 2006;24(3):516–523.
- Shultz SJ, Schmitz RJ. Effects of transverse and frontal plane knee laxity on hip and knee neuromechanics during drop landings. *Am J Sports Med.* 2009;37(9):1821–1830.
- 154. Shultz SJ, Schmitz RJ, Nguyen A, Levine BJ. Joint laxity is related to lower extremity energetics during a drop jump landing. *Med Sci Sports Exerc*. 2010;42(4):771–780.
- 155. Shultz SJ, Schmitz RJ, Kong Y, et al. Cyclic variations in multiplanar knee laxity influence landing biomechanics. *Med Sci Sports Exerc.* 2012;44(5):900–909.
- 156. Shultz SJ, Schmitz RJ, Cone JR, et al. Changes in fatigue, multiplanar knee laxity, and landing biomechanics during intermittent exercise. *J Athl Train*. 2015;50(5):486–497.
- 157. Geiser CF, Meinerz CM, Malloy PJ, Kipp K. Generalized joint hypermobility alters frontal plane knee loading during a drop jump task in Division I female athletes [abstract]. J Athl Train. 2015; 50(10):1104–1105.
- 158. Park SK, Stefanyshyn DJ, Ramage H, Hart DA, Ronsky JL. The relationship between knee joint laxity and knee joint mechanics during the menstrual cycle. *Br J Sports Med.* 2009;43(3):174–179.
- 159. Park SK, Stefanyshyn DJ, Ramage B, Hart DA, Ronsky JL. Alterations in knee joint laxity during the menstrual cycle in healthy women leads to increases in joint loads during selected athletic movements. *Am J Sports Med.* 2009;37(6):1169–1177.
- Shultz SJ, Schmitz RJ, Nguyen A, et al. Knee laxity and its cyclic variations influence tibiofemoral joint motion during weight acceptance. *Med Sci Sports Exerc.* 2011;43(2):287–295.
- Padua DA. Executing a collaborative prospective risk-factor study: findings, successes, and challenges. J Athl Train. 2010;45(5):519– 521.
- 162. Shultz SJ, Nguyen A, Schmitz RJ. Differences in lower extremity anatomical and postural characteristics in males and females between maturation groups. *J Orthop Sports Phys Ther.* 2008; 38(3):137–149.
- Hertel JN, Dorfman JH, Braham RA. Lower extremity malalignments and anterior cruciate ligament injury history. J Sports Sci Med. 2004;3(4):220–225.
- Astrom M, Arvidson T. Alignment and joint motion in the normal foot. J Orthop Sports Phys Ther. 1995;22(5):216–222.
- Loomba-Albrecht LA, Styne DM. Effect of puberty on body composition. *Curr Opin Endocrinol Diabetes Obes*. 2009;16(1):10– 15.
- 166. Montgomery MM, Shultz SJ, Schmitz RJ. The effect of equalizing landing task demands on sex differences in lower extremity energy absorption. *Clin Biomech (Bristol, Avon)*. 2014;29(7):760–766.
- 167. Kulas AS, Hortobagyi T, Devita P. The interaction of trunk-load and trunk-position adaptations on knee anterior shear and hamstrings muscle forces during landing. *J Athl Train.* 2010; 45(1):5–15.
- 168. Taylor JB, Wang H, Schmitz RJ, Rhea CK, Ross SE, Shultz SJ. The influence of multi-planar knee laxity on perceived function during activities of daily living and sport. *J Athl Train*. In press.
- 169. Ficek K, Cieszczyk P, Kaczmarczyk M, et al. Gene variants within the COL1A1 gene are associated with reduced anterior cruciate

ligament injury in professional soccer players. J Sci Med Sport. 2013;16(5):396–400.

- 170. Khoschnau S, Melhus H, Jacobson A, et al. Type I collagen alpha-1 SP1 plymorphism and the risk of cruciate ligament ruptures or shoulder dislocation. *Am J Sports Med.* 2008;36(12):2432–2436.
- 171. O'Connell K, Knight H, Ficek K, et al. Interactions between collagen gene variants and risk of anterior cruciate ligament rupture. *Eur J Sport Sci.* 2015;15(4):341–350.
- Posthumus M, September AV, Keegan M, et al. Genetic risk factors for anterior cruciate ligament ruptures: COL1A1 gene variant. *Br J Sports Med.* 2009;43(5):353–356.
- 173. Posthumus M, September AV, O'Cuinneagain D, van der Merwe W, Schwellnus MP, Collins M. The COL5A1 gene is associated with increased risk of anterior cruciate ligament ruptures in female participants. *Am J Sports Med.* 2009;37(11):2234–2240.
- 174. Posthumus M, September AV, O'Cuinneagain D, van der Merwe W, Schwellnus MP, Collins M. The association between the COL12A1 gene and anterior cruciate ligament ruptures. *Br J Sports Med.* 2010;44(16):1160–1165.
- 175. Stepien-Slodkowska M, Ficek K, Eider J, et al. The +1245g/t polymorphisms in the collagen type I alpha 1 (colla1) gene in Polish skiers with anterior cruciate ligament injury. *Biol Sport*. 2013;30(1):57–60.
- 176. Mannion S, Mtintsilana A, Posthumus M, et al. Genes encoding proteoglycans are associated with the risk of anterior cruciate ligament ruptures. *Br J Sports Med.* 2014;48(22):1640–1646.
- 177. Ficek K, Stepien-Slodkowska M, Kaczmarczyk M, et al. Does the A9285G polymorphism in collagen type XII alpha1 gene associate with the risk of anterior cruciate ligament ruptures? *Balkan J Med Genet*. 2014;17(1):41–46.
- 178. Rahim M, Gibbon A, Hobbs H, et al. The association of genes involved in the angiogenesis-associated signaling pathway with risk of anterior cruciate ligament rupture. *J Orthop Res.* 2014;32(12): 1612–1618.
- 179. Collins M, Rahim M, Mannion S, et al. Investigation of genes involved in the cell signaling pathway with risk of anterior cruciate ligament ruptures [abstract]. *J Athl Train*. 2015;50(10):1104.
- 180. Malila S, Yuktanandana P, Saowaprut S, Jiamjarasrangsi W, Honsawek S. Association between matrix metalloproteinase-3 polymorphism and anterior cruciate ligament ruptures. *Genet Mol Res.* 2011;10(4):4158–4165.
- 181. Posthumus M, Collins M, Van Der Merwe L, et al. Matrix metalloproteinase genes on chromosome 11q22 and the risk of anterior cruciate ligament (ACL) rupture. *Scand J Med Sci Sports*. 2012;22(4):523–533.
- Ribbans WJ, Collins M. Pathology of the tendo Achillis: do our genes contribute? *Bone Joint J.* 2013;95B(3):305–313.
- 183. Baird AE, Carter SD, Innes JF, Ollier WE, Short AD. Genetic basis of cranial cruciate ligament rupture (CCLR) in dogs. *Connect Tissue Res.* 2014;55(4):275–281.
- 184. Baird AE, Carter SD, Innes JF, Ollier W, Short A. Genome-wide association study identifies genomic regions of association for cruciate ligament rupture in Newfoundland dogs. *Anim Genet*. 2014;45(4):542–549.
- 185. John R, Dhillon MS, Prabhakar S, Anand A. Association of gene polymorphisms in COL1A1 and COL12A1 with ACL tears: a study in the Indian population [abstract]. *J Athl Train*. 2015;50(10):1103– 1104.
- 186. Dragoo JL, Lee RS, Benhaim P, Finerman GAM, Hame SL. Relaxin receptors in the human female anterior cruciate ligament. Am J Sports Med. 2003;31(4):577–584.
- 187. Faryniarz DA, Bhargava M, Lajam C, Attia ET, Hannafin JA. Quantitation of estrogen receptors and relaxin binding in human

anterior cruciate ligament fibroblasts. In Vitro Cell Dev Biol Anim. 2006;42(7):176–181.

- 188. Hamlet WP, Liu SH, Panossian V, Finerman GA. Primary immunolocalization of androgen target cells in the human anterior cruciate ligament. J Orthop Res. 1997;15(5):657–663.
- Liu SH, Al-Shaikh RA, Panossian V, Finerman GM. Primary immunolocalization of estrogen and progesterone target cells in the human anterior cruciate ligament. *J Orthop Res.* 1996;14(4):526– 533.
- 190. Lovering RM, Romani WA. Effect of testosterone on the female anterior cruciate ligament. *Am J Physiol Regul Integr Comp Physiol*. 2005;289(1):R15–R22.
- 191. Shultz SJ, Wideman L, Montgomery MM, Beasley KN, Nindl BC. Changes in serum collagen markers, IGF-I and knee joint laxity across the menstrual cycle. *J Orthop Res.* 2012;30(9):1405–1412.
- 192. Casey E, Hameed F, Dhaher YY. The muscle stretch reflex throughout the menstrual cycle. *Med Sci Sports Exerc*. 2014;46(3): 600–609.
- 193. Arendt EA, Bershadsky B, Agel J. Periodicity of noncontact anterior cruciate ligament injuries during the menstrual cycle. J Gend Specif Med. 2002;5(2):19–26.
- 194. Beynnon BD, Johnson RJ, Braun S, et al. The relationship between menstrual cycle phase and anterior cruciate ligament injury: a casecontrol study of recreational alpine skiers. *Am J Sports Med.* 2006; 34(5):757–764.
- 195. Myklebust G, Engebretsen L, Braekken IH, Skjolberg A, Olsen OE, Bahr R. Prevention of anterior cruciate ligament injuries in female team handball players: a prospective intervention study over three seasons. *Clin J Sport Med.* 2003;13(2):71–78.
- 196. Slauterbeck JR, Fuzie SF, Smith MP, et al. The menstrual cycle, sex hormones, and anterior cruciate ligament injury. *J Athl Train*. 2002; 37(3):275–280.
- 197. Wojtys EM, Huston L, Boynton MD, Spindler KP, Lindenfeld TN. The effect of menstrual cycle on anterior cruciate ligament in women as determined by hormone levels. *Am J Sports Med.* 2002; 30(2):182–188.
- 198. Romani W, Patrie J, Curl LA, Flaws JA. The correlations between estradiol, estrone, estriol, progesterone, and sex hormone-binding globulin and anterior cruciate ligament stiffness in healthy, active females. *J Womens Health (Larchmt)*. 2003; 12(3):287–297.
- Slauterbeck JR, Hickox MS, Beynnon BD, Hardy DM. Anterior cruciate ligament biology and its relationship to injury forces. *Orthop Clin North Am.* 2006;37(4):585–591.
- 200. Yu WD, Panossian V, Hatch JD, Liu SH, Finerman GA. Combined effects of estrogen and progesterone on the anterior cruciate ligament. *Clin Orthop Relat Res.* 2001;383:268–281.
- 201. Dehghan F, Muniandy S, Yusof A, Salley N. Sex-steroid regulation of relaxin receptor isoforms (RXFP1 & RXFP2) expression in the patellar tendon and lateral collateral ligament of female WKY rats. *Int J Med Sci.* 2014;11(2):180–191.
- 202. Dehghan F, Muniandy S, Yusof A, Salley N. Testosterone reduces knee passive range of motion and expression of relaxin receptor isoforms via 5α-dihydrotestosterone and androgen receptor binding. *Int J Mol Sci.* 2014;15(3):4619–4634.
- 203. Hashem G, Zhang Q, Hayami T, Chen J, Wang W, Kapila S. Relaxin and B-estradiol modulate targeted matrix degradation in specific synovial joint fibrocartilages: progesterone prevents matrix loss. *Arthritis Res Ther.* 2006;8(4):R98.
- 204. Naqvi T, Duong TT, Hashem G, Shiga M, Zhang Q, Kapila S. MMP induction by relaxin causes cartilage matrix degradation in target synovial joints: receptor profiles correlate with matrix turnover. *Arthritis Res Ther.* 2005;7(1):R1–R11.

- 205. Comerford EJ, Tarlton JF, Avery NC, Bailey AJ, Innes JF. Distal femoral intercondylar notch dimensions and their relationship to composition and metabolism of the canine anterior cruciate ligament. *Osteoarthritis Cartilage*. 2006;14(3):273–278.
- 206. Lee C, Liu X, Smith CL, et al. The combined regulation of estrogen and cyclic tension on fibroblast biosynthesis derived from anterior cruciate ligament. *Matrix Biol.* 2004;23(5):323–329.
- 207. Dragoo JL, Castillo TN, Braun HJ, Ridley BA, Kennedy AC, Golish SR. Prospective correlation between serum relaxin concentration and anterior cruciate ligament tears among elite collegiate female athletes. *Am J Sports Med.* 2011;39(10):2175–2180.
- Takano M, Yamaguchi M, Nakajima R, Fujita S, Kojima T, Kasai K. Effects of relaxin on collagen type I released by stretched human periodontal ligament cells. *Orthod Craniofac Res.* 2009;12(4):282–288.
- 209. Unemori EN, Amento EP. Relaxin modulates synthesis and secretion of procollagenase and collagen by human dermal fibroblasts. *J Biol Chem.* 1990;265(18):10681–10685.
- 210. Kang Y, Choi Y, Yun C, et al. Down-regulation of collagen synthesis and matrix metalloproteinase expression in myofibroblasts from Dupuytren nodule using adenovirus-mediated relaxin gene therapy. *J Orthop Res.* 2014;32(4):515–523.
- Wood ML, Luthin WN, Lester GE, Dahners LE. Tendon creep is potentiated by NKISK and relaxin which produce collagen fiber sliding. *Iowa Orthop J.* 2003;23:75–79.
- Unemori EN, Beck S, Lee WP, et al. Human relaxin decreases collagen accumulation in vivo in two rodent models of fibrosis. J Invest Dermatol. 1993;101(3):280–285.
- Dragoo JL, Padrez K, Workman R, Lindsey DP. The effect of relaxin on the female anterior cruciate ligament: analysis of mechanical properties in an animal model. *Knee*. 2009;16(1):69– 72.
- Wideman L, Montgomery MM, Levine BJ, Beynnon BD, Shultz SJ. Accuracy of calendar-based methods for assigning menstrual cycle phase in women. *Sports Health.* 2013;5(2):143–149.
- 215. Shultz SJ, Wideman L, Montgomery MM, Levine BJ. Some sex hormone profiles are consistent over time in normal menstruating females: implications for sports injury epidemiology. *Br J Sports Med.* 2011;45(9):735–742.
- 216. Vescovi JD. The menstrual cycle and anterior cruciate ligament injury risk. *Sports Med.* 2011;41(2):91–101.
- Bryant GD, Panter ME, Stelmasiak T. Immunoreactive relaxin in human serum during the menstrual cycle. *J Clin Endocrinol Metab.* 1975;41(6):1065–1069.
- 218. Johnson MR, Carter G, Grint C, Lightman SL. Relationship between ovarian steroids, gonadotropins and relaxin during the menstrual cycle. *Acta Endocrinol*. 1993;129(2):121–125.
- Stewart DR, Celniker AC, Taylor CA, Cragun JR, Overstreet JW, Lasley BL. Relaxin in the peri-implantation period. J Clin Endocrinol Metab. 1990;70(6):1771–1773.
- 220. Wreje U, Kristiansson P, Aberg H, Bystrom B, Schoultz B. Serum levels of relaxin during the menstrual cycle and oral contraceptive use. *Gynecol Obstet Invest*. 1995;39(3):197–200.
- 221. Dragoo JL, Castillo TN, Korotkova TA, Kennedy AC, Kim HJ, Stewart DR. Trends in serum relaxin concentration among elite collegiate female athletes. *Int J Womens Health*. 2011;3:19–24.
- 222. Wolf JM, Williams AE, Delaronde S, Leger R, Clifton KB, King KB. Relationship of serum relaxin to generalized and trapezialmetacarpal joint laxity. *J Hand Surg Am.* 2013;38(4):721–728.
- 223. Wolf JM, Cameron KL, Clifton KB, Owens BD. Serum relaxin levels in young athletic men are comparable with those in women. *Orthopaedics*. 2013;36(2):128–131.

- 224. Wolf JM, Scher DL, Etchill EW, et al. Relationahip of relaxin hormone and thumb carpometacarpal joint arthritis. *Clin Orthop Relat Res.* 2014;472(4):1130–1137.
- 225. Pehrsson M, Westberg L, Landen M, Ekman A. Stable serum levels of relaxin throughout the menstrual cycle: a preliminary comparison of women with premenstrual dysphoria and controls. *Arch Womens Ment Health.* 2007;10(4):147–153.
- 226. De Souza MJ, Toombs RJ, Scheid JL, O'Donnell E, West SL, Williams NI. High prevalence of subtle and severe menstrual disturbances in exercising women: confirmation using daily hormone measures. *Hum Reprod*. 2010;25(2):491–503.
- 227. Tourville T, Shultz SJ, Vacek P, et al. Evaluation of an algorithm to predict menstrual cycle phase at the time of injury. *J Athl Train*. In press.
- 228. Burrows M, Peters CE. The influence of oral contraceptives on athletic performance in female athletes. *Sports Med.* 2007;37(7): 557–574.
- 229. Massaro M, Di Carlo C, Gargano V, Formisano C, Bifulco G, Nappi C. Effects of the contraceptive patch and the vaginal ring on bone metabolism and bone mineral density: a prospective, controlled, randomized study. *Contraception*. 2010;881(3):209–214.
- van Mechelen W, Hlobil H, Kemper HC. Incidence, severity, aetiology and prevention of sports injuries: a review of concepts. *Sports Med.* 1992;14(2):82–99.
- 231. Myer GD, Ford KR, Khoury J, Succop P, Hewett TE. Development and validation of a clinic-based prediction tool to identify female athletes at high risk for anterior cruciate ligament injury. *Am J Sports Med.* 2010;38(10):2025–2033.
- 232. Myer GD, Ford KR, Khoury J, Succop P, Hewett TE. Biomechanics laboratory-based prediction algorithm to identify female athletes with high knee loads that increase risk of ACL injury. *Br J Sports Med.* 2011;45(4):245–252.
- 233. Padua DA, Marshall SW, Boling MC, Thigpen CA, Garrett WE, Beutler AI. The Landing Error Scoring System (LESS) is a valid and reliable clinical assessment tool of jump-landing biomechanics: the JUMP-ACL study. *Am J Sports Med.* 2009;37(10):1996–2002.
- Herrington L, Myer GD, Munro A. Intra and inter-tester reliability of the tuck jump assessment. *Phys Ther Sport*. 2013;14(3):152–155.
- 235. DiStefano LJ, Padua DA, DiStefano MJ, Marshall SW. Influence of age, sex, technique, and exercise program on movement patterns after an anterior cruciate ligament injury prevention program in youth soccer players. *Am J Sports Med.* 2009;37(3):495–505.
- 236. Padua DA, DiStefano LJ, Marshall SW, Beutler AI, de la Motte SJ, DiStefano MJ. Retention of movement pattern changes after a lower extremity injury prevention program is affected by program duration. *Am J Sports Med.* 2012;40(2):300–306.
- 237. Hootman JM, Dick R, Agel J. Epidemiology of collegiate injuries for 15 sports: summary and recommendations for injury prevention initiatives. J Athl Train. 2007;42(2):311–319.
- 238. Dragoo JL, Braun HJ, Durham JL, Chen MR, Harris AH. Incidence and risk factors for injuries to the anterior cruciate ligament in National Collegiate Athletic Association football: data from the 2004–2005 through 2008–2009 National Collegiate Athletic Association Injury Surveillance System. *Am J Sports Med.* 2012; 40(5):990–995.
- 239. Shelbourne KD, Gray T, Haro M. Incidence of subsequent injury to either knee within 5 years after anterior cruciate ligament reconstruction with patellar tendon autograft. *Am J Sports Med.* 2009;37(2):246–251.
- Salmon L, Russell V, Musgrove T, Pinczewski L, Refshauge K. Incidence and risk factors for graft rupture and contralateral rupture after anterior cruciate ligament reconstruction. *Arthroscopy*. 2005; 21(8):948–957.

- 241. Paterno MV, Rauh MJ, Schmitt LC, Ford KR, Myer GD, Hewett TE. Incidence of contralateral and ipsilateral anterior cruciate ligament (ACL) injury after primary ACL reconstruction and return to sport. *Clin J Sport Med.* 2012;22(2):116–121.
- 242. Paterno MV, Rauh MJ, Schmitt LC, Ford KR, Hewett TE. Incidence of second ACL injuries 2 years after primary ACL reconstruction and return to sport. *Am J Sports Med.* 2014;42(7):1567–1573.
- 243. Pinczewski LA, Lyman J, Salmon LJ, Russell VJ, Roe J, Linklater J. A 10-year comparison of anterior cruciate ligament reconstructions with hamstring tendon and patellar tendon autograft: a controlled, prospective trial. *Am J Sports Med.* 2007;35(4):564–574.
- 244. Leys T, Salmon L, Waller A, Linklater J, Pinczewski L. Clinical results and risk factors for reinjury 15 years after anterior cruciate ligament reconstruction: a prospective study of hamstring and patellar tendon grafts. *Am J Sports Med.* 2012;40(3):595–605.
- 245. Kaeding CC, Pedroza AD, Reinke EK, Huston LJ, MOON Consortium, Spindler KP. Risk factors and predictors of subsequent ACL Injury in either knee after ACL reconstruction: prospective analysis of 2488 primary ACL reconstructions from the MOON cohort. *Am J Sports Med.* 2015;43(7):1583–1590.
- 246. Myer GD, Ford KR, Khoury J, Hewett TE. Three-dimensional motion analysis validation of a clinic-based nomogram designed to identify high ACL injury risk in female athletes. *Phys Sportsmed*. 2011;39(1):19–28.
- 247. Myer GD, Ford KR, Hewett TE. New method to identify athletes at high risk of ACL injury using clinic-based measurements and freeware computer analysis. *Br J Sports Med*. 2011;45(4):238–244.
- 248. Swart E, Redler L, Fabricant PD, Mandelbaum BR, Ahmad CS, Wang YC. Prevention and screening programs for anterior cruciate ligament injuries in young athletes: a cost-effectiveness analysis. J Bone Joint Surg Am. 2014;96(9):705–711.
- 249. McCall A, Carling C, Nedelec M, et al. Risk factors, testing and preventative strategies for non-contact injuries in professional football: current perceptions and practices of 44 teams from various premier leagues. *Br J Sports Med.* 2014;48(18):1352–1357.
- 250. Gilchrist J, Mandelbaum BR, Melancon H, et al. A randomized controlled trial to prevent noncontact anterior cruciate ligament injury in female collegiate soccer players. *Am J Sports Med.* 2008; 36(8):1476–1483.
- Hewett TE. The effect of neuromuscular training on the incidence of knee injury in female athletes: a prospective study. *Am J Sports Med.* 1999;27(6):699–705.
- 252. Hewett TE, Ford KR, Myer GD. Anterior cruciate ligament injuries in female athletes: part 2, a meta-analysis of neuromuscular interventions aimed at injury prevention. *Am J Sports Med.* 2006; 34(3):1–9.
- 253. Mandlebaum BR, Silvers HJ, Watanabe DS, et al. Effectiveness of a neuromuscular and proprioceptive training program in preventing anterior cruciate ligament injuries in female athletes. *Am J Sports Med.* 2005;33(7):1003–1010.
- 254. Olsen OE, Myklebust G, Engebretsen L, Holme I, Bahr R. Exercises to prevent lower limb injuries in youth sports: cluster randomised controlled trial. *BMJ*. 2005;330(7489):449.
- 255. Yoo JH, Lim BO, Ha M, et al. A meta-analysis of the effect of neuromuscular training on the prevention of the anterior cruciate ligament injury in female athletes. *Knee Surg Sports Traumatol Arthrosc.* 2010;18(6):824–830.
- 256. Sadoghi P, von Keudell A, Vavken P. Effectiveness of anterior cruciate ligament injury prevention training programs. *J Bone Joint Surg Am.* 2012;94(9):769–776.
- 257. LaBella CR, Huxford MR, Grissom J, Kim KY, Peng J, Christoffel KK. Effect of neuromuscular warm-up on injuries in female soccer and basketball athletes in urban public high schools: cluster

randomized controlled trial. Arch Pediatr Adolesc Med. 2011; 165(11):1033–1040.

- 258. Herman DC, Onate JA, Weinhold PS, et al. The effects of feedback with and without strength training on lower extremity biomechanics. *Am J Sports Med.* 2009;37(7):1301–1308.
- 259. Hertel JN, Guskiewicz KM, Kahler DM, Perrin DH. Effect of lateral ankle joint anesthesia on center of balance, postural sway, and joint position sense. *J Sport Rehabil.* 1996;5(2):111–119.
- 260. Hurd WJ, Chmielewski TL, Snyder-Mackler L. Perturbationenhanced neuromuscular training alters muscle activity in female athletes. *Knee Surg Sports Traumatol Arthrosc.* 2006;14(1):60–69.
- Myer GD, Ford KR, Brent JL, Hewett TE. The effects of plyometric vs. dynamic stabilization and balance training on power, balance, and landing force in female athletes. *J Strength Cond Res.* 2006; 20(2):345–353.
- 262. Myer GD, Ford KR, McLean SG, Hewett TE. The effects of plyometric versus dynamic stabilization and balance training on lower extremity biomechanics. *Am J Sports Med.* 2006;34(3):445–455.
- 263. Oñate JA, Guskiewicz KM, Marshall SW, Giuliani C, Yu B, Garrett WE. Instruction of jump-landing technique using videotape feedback: altering lower extremity motion patterns. *Am J Sports Med.* 2005;33(6):831–842.
- 264. Pollard CD, Sigward SM, Ota S, Langford K, Powers CM. The influence of in-season injury prevention training on lower-extremity kinematics during landing in female soccer players. *Clin J Sport Med.* 2006;16(3):223–227.
- 265. Ter Stege MH, Dallinga JM, Benjaminse A, Lemmink KA. Effect of interventions on potential, modifiable risk factors for knee injury in team ball sports: a systematic review. *Sports Med.* 2014;44(10): 1403–1426.
- 266. Leppanen M, Aaltonen S, Parkkari J, Heinonen A, Kujala UM. Interventions to prevent sports related injuries: a systematic review and meta-analysis of randomised controlled trials. *Sports Med.* 2014;44(4):473–486.
- 267. Stevenson JH, Beattie CS, Schwartz JB, Busconi BD. Assessing the effectiveness of neuromuscular training programs in reducing the incidence of anterior cruciate ligament injuries in female athletes: a systematic review. *Am J Sports Med.* 2015;43(2):482–490.
- 268. Hewett TE, Stroupe AL, Noyes FR. Plyometric training in female athletes: decreased impact forces and increased hamstring torques. *Am J Sports Med.* 1996;24(6):765–773.
- Alentorn-Geli E, Mendiguchia J, Samuelsson K, et al. Prevention of anterior cruciate ligament injuries in sports: part I, systematic review of risk factors in male athletes. *Knee Surg Sports Traumatol Arthrosc.* 2014;22(1):3–15.
- 270. Sugimoto D, Alentorn-Geli E, Mendiguchia J, Samuelsson K, Karlsson J, Myer GD. Biomechanical and neuromuscular characteristics of male athletes: implications for the development of anterior cruciate ligament injury prevention programs. *Sports Med.* 2015;45(6):809–822.
- Wulf G, Chiviacowsky S, Schiller E, Avila LT. Frequent externalfocus feedback enhances motor learning. *Front Psychol.* 2010;1:1– 7.
- 272. Pappas E, Nightingale EJ, Simic M, Ford KR, Hewett TE, Myer GD. Do exercises used in injury prevention programmes modify cutting task biomechanics? A systematic review with meta-analysis. *Br J Sports Med.* 2015;49(10):673–680.
- 273. Benjaminse A, Otten B, Gokeler A, Diercks RL, Lemmink KAPM. Sex specific motor learning strategies: implications for ACL injury prevention [abstract]. *J Athl Train*. 2015;50(10):1110–1111.
- 274. Gokeler A, Welling W, Otten B, Benjaminse A. Enhanced retention of drop vertical jump landing strategies assessed with the Landing Error Scoring System [abstract]. J Athl Train. 2015;50(10):1110.

- 275. Dowling AV, Favre J, Andriacchi TP. Inertial sensor-based feedback can reduce key risk metrics for anterior cruciate ligament injury during jump landings. *Am J Sports Med.* 2012;40(5):1075–1083.
- 276. Barrios JA, Crossley KM, Davis IS. Gait retraining to reduce the knee adduction moment through real-time visual feedback of dynamic knee alignment. *J Biomechanics*. 2010;43(11):2208–2213.
- 277. Crowell HP, Milner CE, Hamill J, Davis IS. Reducing impact loading during running with the use of real-time visual feedback. J Orthop Sports Phys Ther. 2010;40(4):206–213.
- 278. Ford KR, DiCesare CA, Myer GD, Hewett TE. Real-time biofeedback to target risk of anterior cruciate ligament injury: a technical report for injury prevention and rehabilitation. *J Sport Rehabil.* 2015; Technical Report 13. doi:10.1123/jsr.2013-0138.
- 279. Munro A, Herrington L. The effect of videotape augmented feedback on drop jump landing strategy: implications for anterior cruciate ligament and patellofemoral joint injury prevention. *Knee*. 2014;21(5):891–895.
- 280. Myer GD, Stroube BW, DiCesare CA, et al. Augmented feedback supports skill transfer and reduces high-risk injury landing mechanics: a double-blind, randomized controlled laboratory study. *Am J Sports Med.* 2013;41(3):669–677.
- 281. Stroube BW, Myer GD, Brent JL, Ford KR, Heidt RS Jr, Hewett TE. Effects of task-specific augmented feedback on deficit modification during performance of the tuck-jump exercise. *J Sport Rehabil.* 2013;22(1):7–18.
- 282. Benjaminse A, Gokeler A, Dowling AV, et al. Optimization of the anterior cruciate ligament injury prevention paradigm: novel feedback techniques to enhance motor learning and reduce injury risk. J Orthop Sports Phys Ther. 2015;45(3):170–182.
- 283. DiStefano LJ, Blackburn JT, Marshall SW, Guskiewicz KM, Garrett WE, Padua DA. Effects of an age-specific anterior cruciate ligament injury prevention program on lower extremity biomechanics in children. *Am J Sports Med.* 2011;39(5):949–957.
- 284. Frank BS, Register-Mihalik J, Padua DA. High levels of coach intent to integrate a ACL injury prevention program into training does not translate to effective implementation. J Sci Med Sport. 2015;18(4):400–406.
- 285. Sugimoto D, Myer GD, Bush HM, Klugman MF, Medina McKeon JM, Hewett TE. Compliance with neuromuscular training and anterior cruciate ligament injury risk reduction in female athletes: a meta-analysis. *J Athl Train*. 2012;47(6):714–723.
- 286. McKay CD, Steffen K, Romiti M, Finch CF, Emery CA. The effect of coach and player injury knowledge, attitudes and beliefs on adherence to the FIFA 11+ programme in female youth soccer. *Br J Sports Med.* 2014;48(17):1281–1286.
- 287. Lindblom H, Walden M, Carlfjord S, Hagglund M. Implementation of a neuromuscular training programme in female adolescent football: 3-year follow-up study after a randomised controlled trial. *Br J Sports Med.* 2014;48(19):1425–1430.
- 288. Joy EA, Taylor JR, Novak MA, Chen M, Fink BP, Porucznik CA. Factors influencing the implementation of anterior cruciate ligament injury prevention strategies by girls soccer coaches. *J Strength Cond Res.* 2013;27(8):2263–2269.
- 289. Norcross MF, Johnson ST, Bovbjerg VE, Koester MC, Hoffman MA. Factors influencing high school coaches' adoption of injury prevention programs. J Sci Med Sport. In press.
- Ekstrand J. Keeping your top players on the pitch: the key to football medicine at a professional level. *Br J Sports Med.* 2013; 47(12):723–724.
- 291. Martinez JC, Roux E, Eason CM, et al. Factors that influence female adolescent athletes' willingness to perform an injury prevention program [abstract]. *J Athl Train*. 2015;50(10):1112.

- 292. Keats MR, Emery CA, Finch CF. Are we having fun yet? Fostering adherence to injury preventive exercise recommendations in young athletes. *Sports Med.* 2012;42(3):175–184.
- 293. Myer GD, Ford KR, Palumbo JP, Hewett TE. Neuromuscular training improves performance and lower-extremity biomechanics in female athletes. *J Strength Cond Res.* 2005;19(1):51–60.
- 294. Benjaminse A, Welling W, Otten B, Gokeler A. Novel methods of instruction in ACL injury prevention programs, a systematic review. *Phys Ther Sport.* 2015;16(2):176–186.
- 295. Padua DA, Frank B, Donaldson A, et al. Seven steps for developing and implementing a preventive training program: lessons learned from JUMP-ACL and beyond. *Clin Sports Med.* 2014;33(4):615– 632.
- 296. O'Brien J, Finch CF. The implementation of musculoskeletal injury-prevention exercise programmes in team ball sports: a systematic review employing the RE-AIM framework. *Sports Med.* 2014;44(9):1305–1318.
- 297. Sugimoto D, Myer GD, Foss KD, Hewett TE. Specific exercise effects of preventive neuromuscular training intervention on anterior cruciate ligament injury risk reduction in young females: meta-analysis and subgroup analysis. *Br J Sports Med.* 2015;49(5): 282–289.
- 298. Myer GD, Sugimoto D, Thomas S, Hewett TE. The influence of age on the effectiveness of neuromuscular training to reduce anterior cruciate ligament injury in female athletes: a meta-analysis. *Am J Sports Med.* 2013;41(1):203–215.
- 299. Taylor JB, Waxman JP, Richter SJ, Shultz SJ. Evaluation of the effectiveness of anterior cruciate ligament injury prevention programme training components: a systematic review and metaanalysis. *Br J Sports Med.* 2015;49(2):79–87.
- 300. Palmieri-Smith RM, Wojtys EM, Ashton-Miller JA. Association between preparatory muscle activation and peak valgus knee angle. *J Electromyogr Kinesiol*. 2008;18(6):973–979.
- 301. Pollard CD, Sigward SM, Powers CM. Limited hip and knee flexion during landing is associated with increased frontal plane knee motion and moments. *Clin Biomech (Bristol, Avon)*. 2010;25(2): 142–146.
- 302. Claiborne TL, Armstrong CW, Gandhi V, Pincivero DM. Relationship between hip and knee strength and knee valgus during a single leg squat. J Appl Biomech. 2006;22(6):41–50.
- 303. Russell KA, Palmieri RM, Zinder SM, Ingersoll CD. Sex differences in valgus knee angle during a single-leg drop jump. J Athl Train. 2006;41(2):166–171.
- 304. Zazulak BT, Ponce PL, Straub SJ, Medvecky MJ, Avedisian L, Hewett TE. Gender comparison of hip muscle activity during single-leg landing. J Orthop Sports Phys Ther. 2005;35(5):292–299.
- 305. Hollman JH, Ginos BE, Kozuchowski J, Vaughn AS, Krause DA, Youdas JW. Relationships between knee valgus, hip-muscle strength, and hip-muscle recruitment during a single-limb stepdown. J Sport Rehabil. 2009;18(1):104–117.
- 306. Lawrence RK, Kernozek TW, Miller EJ, Torry MR, Reuteman P. Influences of hip external rotation strength on knee mechanics during single-leg drop landings in females. *Clin Biomech (Bristol, Avon)*. 2008;23(6):806–813.
- 307. Ford KR, Myer GD, Hewett TE. Longitudinal effects of maturation on lower extremity joint stiffness in adolescent athletes. *Am J Sports Med.* 2010;38(9):1829–1837.
- 308. Carcia CR, Kivlan B, Scibek JS. The relationship between lower extremity closed kinetic chain strength and sagittal plane landing kinematics in female athletes. *Int J Sports Phys Ther.* 2011;6(1):1– 9.
- 309. Mizner RL, Kawaguchi JK, Chmielewski TL. Muscle strength in the lower extremity does not predict postinstruction improvements

in the landing patterns of female athletes. J Orthop Sports Phys Ther. 2008;38(6):353–361.

- 310. McGlashan AJ, Finch CF. The extent to which behavioural and social sciences theories and models are used in sport injury prevention research. *Sports Med.* 2010;40(10):841–858.
- 311. Wulf G, Shea C, Lewthwaite R. Motor skill learning and performance: a review of influential factors. *Med Educ.* 2010; 44(1):75–84.
- 312. Gokeler A, Benjaminse A, Hewett TE, et al. Feedback techniques to target functional deficits following anterior cruciate ligament reconstruction: implications for motor control and reduction of second injury risk. *Sports Med.* 2013;43(11):1065–1074.
- 313. Gokeler A, Benjaminse A, Welling W, Alferink M, Eppinga P, Otten B. The effects of attentional focus on jump performance and

knee joint kinematics in patients after ACL reconstruction. *Phys Ther Sport*. 2015;16(2):114–120.

- 314. Gokeler A, Bisschop M, Myer GD, et al. Immersive virtual reality improves movement patterns in patients after ACL reconstruction: implications for enhanced criteria-based return-to-sport rehabilitation. *Knee Surg Sports Traumatol Arthrosc.* In press.
- Chiviacowsky S, Wulf G. Self-controlled feedback is effective if it is based on the learner's performance. *Res Q Exerc Sport.* 2005; 76(1):42–48.
- 316. Andrieux M, Danna J, Thon B. Self-control of task difficulty during training enhances motor learning of a complex coincidenceanticipation task. *Res Q Exerc Sport.* 2012;83(1):27–35.
- 317. Chiviacowsky S, Wulf G. Self-controlled feedback: does it enhance learning because performers get feedback when they need it? *Res Q Exerc Sport*. 2002;73(4):408–415.

Address correspondence to Sandra J. Shultz, PhD, ATC, FNATA, FACSM, Department of Kinesiology, University of North Carolina at Greensboro, 1408 Walker Avenue, Greensboro, NC 27412. Address e-mail to sjshultz@uncg.edu.