

# Research at the Point of Care: Using Electronic Medical Record Systems to Generate Clinically Meaningful Evidence

Ashley N. Marshall, PhD, ATC\*; Kenneth C. Lam, ScD, ATC†

\*Department of Health and Exercise Science, Appalachian State University, Boone, NC; †Department of Interdisciplinary Health Sciences, A.T. Still University, Mesa, AZ

**Context:** Health care leaders have recommended the use of health information technology to improve the quality of patient care. In athletic training, using informatics, such as electronic medical records (EMRs), would support practice-based decisions about patient care. However, athletic trainers (ATs) may lack the knowledge to effectively participate in point-of-care clinical research using EMRs.

**Objectives:** To discuss the role of EMRs in athletic training and identify methodologic approaches to conducting clinical research at the point of care.

**Description:** The 2020 Commission on Accreditation of Athletic Training Education curricular content standards included the use of an electronic patient record to document care, mitigate error, and support decision making through the collection and use of patient data (Standard 64). Patient data are collected by ATs at the point of care via routine documentation, and these data can be used to answer clinical questions about their practice. Observational or descriptive

study designs are ideal for this type of data. Observational research (ie, case-control, cross-sectional, cohort studies) evaluates factors that influence patients' lives in the "real world," whereas descriptive research (ie, case study or series, descriptive epidemiology studies) identifies characteristics of individuals and groups. If ATs are comprehensively documenting patient care using an EMR, they have the means to participate in observational and descriptive research.

**Clinical and Research Advantages:** Using an EMR to its full capacity allows ATs to collect meaningful data at the point of care, conduct practice-based research, and improve health care for the patient and clinician. However, to ensure data quality, these approaches must include routine and comprehensive documentation habits.

**Key Words:** practice-based research, patient care documentation, quality improvement

The widespread call to leverage health information technology (HIT) for patient care can be traced back 2 decades to the Institute of Medicine's landmark publication, *Crossing the Quality Chasm*.<sup>1</sup> Since then, the National Academy of Medicine has consistently advocated for the implementation of HIT to meet the complex needs of the 21st-century patient,<sup>2</sup> facilitate the exchange of health information,<sup>2,3</sup> deliver high-quality patient-centered care,<sup>2,4,5</sup> and advance research in health care.<sup>2,6,7</sup> Additionally, the US health care system has become increasingly digitized as a result of federal initiatives such as the Health Information Technology for Economic and Clinical Health Act.<sup>8</sup> This act incentivized and supported the implementation of electronic records with the aims of (1) improving quality, safety, and efficiency and reducing health disparities, (2) increasing patient engagement, (3) improving care coordination, (4) expanding population and public health, and (5) ensuring adequate privacy and security protection for personal health.<sup>9</sup>

As the most commonly recognized form of HIT, electronic records are crucial to the patient-clinician relationship and allow information to be "interactive, real time, and prospective."<sup>1</sup> In general, electronic records offer a method for collecting meaningful patient data that support and improve patient care. On a global health care level,

electronic records have been used to better understand disease diagnoses and prognoses,<sup>10,11</sup> improve disease treatment and management strategies,<sup>12,13</sup> reduce health care costs,<sup>14,15</sup> and improve patient outcomes.<sup>16</sup> Because of these benefits, members of the Athletic Training Strategic Alliance<sup>17</sup> have encouraged the use of HIT in athletic training, including electronic records. Although the terms *electronic medical record* (EMR) and *electronic health record* (EHR) are often used interchangeably, they are different. Electronic medical records are standalone systems that reside in a single location and do not interact with other electronic patient records, whereas EHRs have various levels of connectivity between providers or entities so that information can be shared more efficiently and data can be aggregated.<sup>18</sup> Some athletic trainers (ATs) employed by hospitals or health services systems may use an EHR for patient documentation, but most ATs likely use EMRs as part of their clinical practice.

## THE ROLE OF EMRs IN ATHLETIC TRAINING

The Commission on Accreditation of Athletic Training Education<sup>19</sup> recently published the "2020 Standards for Accreditation of Professional Athletic Training Programs." The new curricular content standards include the use of an EMR to document care, mitigate error, and support decision

making through the collection and use of patient data under the health care informatics core competency (Standard 64).<sup>19</sup> This core competency corresponds well<sup>20</sup> with EMR-related recommendations described in the Board of Certification Practice Analysis<sup>21</sup> and the National Athletic Trainers' Association Educational Competencies,<sup>22</sup> which support the usefulness of HIT and EMRs in athletic training. Most recently, the Athletic Training Strategic Alliance's document, "The Prioritized Research Agenda for the Athletic Training Profession,"<sup>17</sup> highlighted the importance of HIT and patient documentation data in advancing athletic training practice and improving the quality of patient care. Taken together, these efforts emphasize the vital role HIT, EMRs, and patient care data will play in the future of our profession.

Although most EMR-related initiatives are aimed at improving "immediate" patient care (ie, patient care documentation, communication, access to patient records), EMRs can be useful for conducting point-of-care clinical research on a short-term or long-term basis if leveraged properly. Collecting patient data at the point of care through routine patient care documentation allows clinicians to deliver effective, evidence-based services and reduce research-specific clinic visits and overall health care costs for the patient.<sup>23</sup> Despite these potential benefits, it is evident that ATs do not comprehensively document care and use EMRs to their full capacity.<sup>24–26</sup> Several groups have analyzed patient care documentation in athletic training, particularly for injury surveillance. For example, researchers have used the National Collegiate Athletic Association Injury Surveillance Program and High School Reporting Information Online system for a decade to describe the injury and exposure data of athletes participating in 13 collegiate and high school sports.<sup>27</sup> Although these data-collection systems yield reliable and valid findings about injury incidence in various sport participation settings, additional opportunities exist to improve care beyond tracking injuries and capturing descriptive data.

## A CALL TO ACTION: USING EMRs TO CONDUCT RESEARCH AT THE POINT OF CARE IN ATHLETIC TRAINING

### Observational Studies

If ATs are comprehensively documenting patient care using commonly recorded variables (Table 1), they should be able to support observational research efforts and use EMR data in a clinically meaningful way. Observational research is performed in a "real-world" clinical setting and evaluates factors that influence patients' lives as they naturally exist.<sup>28</sup> Associations between the recorded variables and outcomes of interest are then analyzed. Although observational studies lack the internal validity of randomized controlled trials, their strong external validity can provide more clinically meaningful findings for athletic health care.<sup>29,30</sup>

Several approaches can be used when conducting observational research using data derived from an EMR (Tables 2 and 3). Because clinicians using an EMR have existing data, retrospective studies are recommended as a first step. One retrospective approach is a case-control study<sup>31</sup> (Figure 1 and Table 2), in which the goal is to establish a relationship between exposures (ie, intervention,

**Table 1. Common Variables Athletic Trainers Document in an Electronic Medical Record System and Common Clinical Questions**

Common Variables	Common Clinical Questions
Sex	What is the most common injury that I see in my practice?
Sport	Are there differences in injury rates based on sport or sex?
Diagnosis (ICD codes)	Is variable X associated with injury Y?
Mechanism of injury	Is treatment X associated with good outcomes?
Symptoms	
Activity level	
Treatment or management (CPT codes)	
Time to return to sport	
Clinician-rated outcome measures	Does the presence of symptom X on evaluation result in a longer recovery time?
Patient-rated outcome measures	

Abbreviations: ICD, International Classification of Diseases; CPT, Current Procedural Terminology.

characteristic, or experience<sup>32</sup>) and an outcome. Another approach is the cross-sectional study design<sup>33</sup> (Figure 2 and Table 2), which compares multiple groups at 1 point in time. The prospective cohort study design<sup>34</sup> (Figure 3 and Table 2) should also be considered; its goal is to determine whether a particular recorded variable is associated with an outcome. However, it is important to note that due to its prospective nature, this design requires an additional level of planning.

### Example of an Observational Study

Teel et al<sup>31</sup> recently investigated the effect of several acute variables on concussion-recovery patterns in collegiate and high school athletes. Variables were collected at baseline (eg, concussion history) or at initial injury evaluation (eg, amnesia, loss of consciousness) by an AT participating in the Concussion Prevention Initiative.<sup>31</sup> Patients were tracked for 90 days postinjury and their data were extracted from the Concussion Prevention Initiative database by the investigators.<sup>31</sup> This is an example of a study that could have been conducted with an EMR. To comply with current recommendations for best practice,<sup>35</sup> ATs should already be documenting such variables in their EMRs. Thus, a case-control study could be conducted that retrospectively relates concussion-evaluation variables to the classification of patients as having experienced prolonged recovery (cases) versus normal (controls) recovery.

### Descriptive Studies


Descriptive studies are another type of clinical research that can be supported by an EMR. The aim of descriptive studies is to examine the behaviors, conditions, and characteristics of individuals and groups.<sup>36</sup> Common approaches to descriptive studies are the case study and series,<sup>37</sup> in which the goal is to depict the characteristics of a patient or group of patients with a rare condition, and descriptive epidemiology designs,<sup>38</sup> in which the goal is to describe injuries, illnesses, or conditions in relation to time, place, and patient (Tables 1 and 2). Studies based on data from the Athletic Training Practice-Based Research Network<sup>26,39–41</sup> and National Athletic Treatment, Injury and Outcomes Network<sup>42,43</sup> have demonstrated the mean-


**Table 2. Study Approaches to Data Derived From an Electronic Medical Record System**

Study Design	Basic Approach	Options for Use of Data	Research Question Example
Case-control	<p>Purpose: To establish a relationship between exposures and specific outcomes</p> <p>Study design moves from outcome to exposure</p> <p>Identify a population with a specific outcome (ie, cases) and select a control group for comparison</p> <p>The control population should not differ in major characteristics from the case population</p> <p>Matching patients between case and control populations limits confounding variables</p>	<p>Compare preseason screening data between those who sustained an injury and those who did not</p> <p>Compare initial injury presentation and symptoms between those who experienced a normal recovery and those who experienced a prolonged recovery</p> <p>Compare the injury-treatment and -management strategies used for those who went on to experience recurrent or subsequent injury and those who did not</p>	Do sideline concussion assessments predict subsequent neurocognitive impairment after sport-related concussion? <sup>31</sup>
Cross-sectional	<p>Purpose: To estimate the prevalence or proportion of an injury or condition and the relationship between variables and an outcome</p> <p>Several subsets of a group studied at 1 time</p> <p>Conclusions drawn about a population after comparing important characteristics of the subset</p>	<p>Compare patient-reported or clinician-rated variables collected at different times after injury or illness</p> <p>Compare injury presentation and symptoms among age groups or between sexes</p> <p>Document the type of surgery used for a particular condition and compare outcome variables among different surgical techniques</p>	How does functional performance differ between females who underwent an anterior cruciate ligament reconstruction with a bone-patellar tendon-bone versus a semitendinosus or gracilis graft? <sup>33</sup>
Prospective cohort	<p>Purpose: To determine whether a particular characteristic is associated with an outcome of interest (eg, injury, illness)</p> <p>Group followed prospectively over time, with measurements repeated at specific intervals</p> <p>Participants selected with respect to particular characteristics</p> <p>Enrolled individuals (with and without the identified characteristic) are followed over time to observe a particular outcome</p>	<p>Collect preseason baseline measures and determine at the end of the season if injury rates differed between those at risk and those not at risk</p> <p>Evaluate if immediate reporting of an injury improves recovery, performance, or long-term outcomes</p> <p>Determine whether participation in or compliance with preventive training programs is related to injury incidence</p>	Which clinical tests best demonstrate the risk of lateral ankle sprain among high school and collegiate football players? <sup>34</sup>
Case series or case study	<p>Purpose: To describe the characteristics of a single patient or a group of patients with a rare injury or condition, those who have had a particular or unique response to treatment or management, or those who have experienced a specific procedure</p> <p>Individuals selected based on the outcome of interest, and the design does not include a control group</p>	<p>Identify a single patient or group of patients with a rare injury, illness, or condition and describe the</p> <ul style="list-style-type: none"> <li>• Presentation</li> <li>• Evaluation</li> <li>• Treatment or management strategies</li> <li>• Patient-reported or clinician-rated (or both) outcomes</li> </ul>	Is a prophylactic protocol successful in managing an athlete with hemophilia who is playing at a high level of contact sports? <sup>37</sup>
Descriptive epidemiology	<p>Purpose: To describe the injuries, illnesses, or conditions that occur with consideration to time, place, and participants</p> <p>Personal characteristics (eg, age, sex, socioeconomic status, use of medications) may affect the occurrence of an injury or condition and are often analyzed independently and in combination</p>	Collect data on exposures and injuries to calculate injury proportions and rates, demographic variables on the participants themselves, and data surrounding the injury (eg, diagnosis, mechanism, site)	What are the injury rates, locations, and mechanisms of injury in collegiate ultimate players? <sup>38</sup>

**Table 3. Study Design Advantages and Disadvantages**

Study Design	Advantages	Disadvantages
Case-control	Efficient for studying rare outcomes Appropriate for studying outcomes with long induction periods Typically less labor intensive and time consuming than cohort studies	Inefficient for studying rare exposures Retrospective nature can lead to increased bias Retrospective nature limits ability to establish causation
Cross-sectional	Used more frequently than the longitudinal approach due to its efficiency Participants are only tested once; no follow-up visits Results are generally applicable to the population that is studied	Difficult to determine to what extent research findings reflect the effects of time versus other sampling variables Difficult to establish causal relationships between exposures and the outcome of interest Only shows associations between exposures and outcome of interest because of the lack of temporality Results are not applicable to populations other than that studied
Prospective cohort	May provide more clinically relevant and generalizable evidence than randomized controlled trials Efficient for studying (1) rare exposures and (2) multiple outcomes associated with a single exposure Can demonstrate a prospective relationship between exposures and outcomes Provides good descriptive information about exposures	Inefficient for studying (1) rare outcomes or (2) outcomes with long induction periods Can be labor intensive and time consuming
Case series or case study	Allows for an understanding of the entirety of an individual's condition and provides a foundation for clinical research and practice Contributes valuable information about patients' experiences and ultimately helps to improve patient care Often provides preliminary data for hypothesis generation and the development of future research studies on relationships and associations	The relationship between an individual or particular group's characteristics and an outcome or variable of interest cannot be determined Selection of patients may be biased
Descriptive epidemiology	Provides insight into the mechanism of injury or condition, factors that increase risk, and other potential contributors Can identify communities, groups of individuals, or patients who are at increased risk	Describes distribution of variables without regard to causal relationships

**Figure 1. Case-control study design.**



**Figure 2. Cross-sectional study design.**


ingfulness of descriptive designs. Using data derived from these networks, researchers have offered insights into athletic training clinical practice, including common injuries managed,<sup>26,39</sup> evaluation techniques employed,<sup>44</sup> treatments and services provided,<sup>26,39–43</sup> and estimated direct costs recorded by ATs.<sup>41</sup>

Descriptive studies are also valuable for quality improvement initiatives, which have been a recent focus in athletic training education<sup>19</sup> and practice.<sup>45,46</sup> The first step of any quality improvement effort is to describe a problem that is based on data collected during patient care. Using the STEEP (safe, timely, effective, efficient, equitable, patient-centered) framework for improvement,<sup>1</sup> ATs can identify

potential practice gaps by analyzing patient data. If a problem is identified, a quality improvement initiative can be developed to address the gap, and the EMR can be used to measure whether any improvements occur.<sup>45</sup>

### Example of a Descriptive Study and Quality Improvement

A simple but meaningful example of how a descriptive study drives quality improvement involves looking at the evaluation patterns of ATs to determine whether best practices are being followed. The Ottawa Ankle Rules<sup>47</sup> are well-known and accepted practice guidelines for referring



**Figure 3. Prospective cohort study design.**



patients for radiographs after a foot or ankle injury. Thus, ATs could retrospectively analyze all patients with foot or ankle injuries who were referred for radiographs to determine whether they actually met the Ottawa Ankle Rules criteria. If the percentage of patients meeting the radiograph and Ottawa Ankle Rules requirements was relatively low (eg, 70% or less), then a continuous quality improvement initiative could be instituted to improve that percentage over time. Potential improvements could include implementing educational strategies for ATs, providing clinical decision support system features within the EMR, and establishing a departmental policy for a successive plan-do-study-act cycle.<sup>48</sup>

## CONSIDERATIONS

An advantage of using an EMR is its ability to facilitate access to meaningful, accurate, reliable, and complete data.<sup>49</sup> However, a limitation of conducting research this way is that the quality of the data depends on the quality of the patient care documentation. Inaccurate, incomplete, or missing health information may adversely affect patient care.<sup>49</sup> As a result, to support EMR-based research efforts, comprehensive and consistent patient care documentation is crucial.

The use of an EMR has been shown to improve patient care documentation practices<sup>50</sup> and communication among providers<sup>51</sup> without increasing the time that clinicians spend documenting.<sup>52</sup> Yet, numerous health care providers (eg, physicians,<sup>53</sup> nurses,<sup>54</sup> pharmacists<sup>55</sup>) often cite barriers to high-quality documentation. Although ATs recognize the importance of patient care documentation, they identified time, uncertainty regarding what to document, and facility and personnel resources as obstacles to documentation.<sup>25</sup> Standard operating procedures and standardized reporting methods may provide strategies for improving documentation habits.<sup>17,56</sup> In addition, comprehensive and continuous EMR training preserves data quality,<sup>56,57</sup> and documentation peer reviews can hold clinicians accountable for their documentation practices.<sup>57</sup> Thus, these strategies should be used before and during EMR-based research efforts to optimize data quality.

The mechanics of patient care documentation also affect the ability to conduct quality research using an EMR. For example, baseline testing<sup>58</sup> or injury registration<sup>59</sup> is often separate from injury documentation, requiring clinicians to document in multiple systems and researchers to compile data from multiple sources to analyze the data prospectively. Because a perceived lack of time is a reported barrier to quality medical record keeping,<sup>25</sup> reducing or eliminating the documentation of information in multiple systems is recommended to improve data quality, reduce data redundancy, and increase the feasibility of EMR-based research.

A potential challenge to EMR-based research efforts in athletic training is the lack of continuity between medical records maintained by ATs and those maintained by the patient's external physician or health care system. Athletic trainers are often outside the formal health care system,<sup>60</sup> so extracting comprehensive data from the electronic patient records of multiple health care providers can be difficult. This may hinder an AT's ability to answer certain research questions. For instance, unless the care provided by

external providers during rehabilitation is considered, studies investigating recovery trajectories may not accurately reflect the effectiveness of certain treatments because data points from external providers will be missing.

## CONCLUSIONS

Over the last 2 decades, the use of HIT has been emphasized in the US health care system. In athletic training, recent efforts have highlighted the vital role EMRs may play in improving the quality of patient care and advancing the profession. Although the use of EMRs has primarily focused on injury surveillance, ATs have a valuable opportunity to collect patient care data in a manner that could drive clinically meaningful research, including observational studies, descriptive studies, and quality improvement projects. These types of studies can ultimately benefit ATs and their patients, but proactive planning is required to ensure that the data collected are of high quality. Implementing strategies such as standard operating procedures, continuous EMR training, peer-to-peer documentation audits, and limiting entry of information into multiple systems can enhance and preserve data quality and increase the feasibility of EMR-based research efforts. Although the athletic training profession has made progress in the use of point-of-care data to support injury-surveillance and practice characterization studies, future investigators should conduct prospective, cost-analysis, and comparative effectiveness research using electronic records.

## REFERENCES

1. Institute of Medicine Committee on Quality of Health Care in America. *Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington, DC: National Academies Press; 2001.
2. Whicher D, Rosengren K, Siddiqi S, Simpson L. *The Future of Health Services Research: Advancing Systems Research and Practice in the United States*. Washington, DC: National Academies Press; 2018.
3. Pronovost P, Johns MME, Palmer S, et al., eds. *Procuring Interoperability: Achieving High-Quality, Connected, and Person-Centered Care*. Washington, DC: National Academy of Medicine; 2018.
4. Dzau VJ, McClellan MB, McGinnis JM, Finkelman EM, eds. *Vital Directions for Health & Health Care: An Initiative of the National Academy of Medicine*. Washington, DC: National Academy of Medicine; 2017.
5. Institute of Medicine. *Best Care at Lower Cost: The Path to Continuously Learning Health Care in America*. Washington, DC: National Academies Press; 2013.
6. Selby J, Larson E, Kaushal R, Zirkle M, Hamilton Lopez M, Whicher D. *Accelerating Medical Evidence Generation and Use: Summary of a Meeting Series*. Washington, DC: National Academy of Medicine; 2017.
7. Institute of Medicine. *Redesigning the Clinical Effectiveness Research Paradigm: Innovation and Practice-Based Approaches: Workshop Summary*. Washington, DC: National Academies Press; 2010.
8. Washington V, DeSalvo K, Mostashari F, Blumenthal D. The HITeCH era and the path forward. *N Engl J Med*. 2017;377(10):904–906.
9. Blumenthal D. Stimulating the adoption of health information technology. *N Engl J Med*. 2009;360(15):1477–1479.

10. Singh H, Hirani K, Kadiyala H, et al. Characteristics and predictors of missed opportunities in lung cancer diagnosis: an electronic health record-based study. *J Clin Oncol*. 2010;28(20):3307–3315.
11. Kinn JW, Marek JC, O'Toole MF, Rowley SM, Bufalino VJ. Effectiveness of the electronic medical record in improving the management of hypertension. *J Clin Hypertens (Greenwich)*. 2008;4(6):415–419.
12. Brindis RG, Fitzgerald S, Anderson HV, Shaw RE, Weintraub WS, Williams JF. The American College of Cardiology National Cardiovascular Data Registry (ACC-NCDR): building a national clinical data repository. *J Am Coll Cardiol*. 2001;37(8):2240–2245.
13. Chung SC, Gedeberg R, Nicholas O, et al. Acute myocardial infarction: a comparison of short-term survival in national outcome registries in Sweden and the UK. *Lancet*. 2014;383(9925):1305–1312.
14. De Moor G, Sundgren M, Kalra D, et al. Using electronic health records for clinical research: the case of the EHR4CR project. *J Biomed Inform*. 2015;53:162–173.
15. Eisenstein EL, Collins R, Cracknell B, et al. Sensible approaches for reducing clinical trial costs. *Clin Trials*. 2008;5(1):75–84.
16. Carberry K, Landman Z, Xie M, Feeley T, Henderson J, Fraser C Jr. Incorporating longitudinal pediatric patient-centered outcome measurement into the clinical workflow using a commercial electronic health record: a step toward increasing value for the patient. *J Am Med Inform Assoc*. 2016;23(1):88–93.
17. Eberman LE, Walker SE, Floyd RT, et al. The prioritized research agenda for the athletic training profession: a report from the Strategic Alliance Research Agenda Task Force. *J Athl Train*. 2019;54(2):237–244.
18. Wager K, Lee F, Glaser J. *Health Care Information Systems: A Practical Approach for Health Care Management*. 4th ed. San Francisco, CA: Jossey-Bass; 2017.
19. 2020 Standards for the accreditation of professional athletic training programs. Commission on Accreditation of Athletic Training Education Web site. <https://caate.net/wp-content/uploads/2018/02/2020-Standards-for-Professional-Programs-copyedited-clean.pdf>. Accessed July 5, 2018.
20. 2020 Standards for the accreditation of professional athletic training programs crosswalk. Commission on Accreditation of Athletic Training Education Web site. [https://caate.net/wp-content/uploads/2018/02/2020-Standards-Crosswalk\\_Final\\_for-Professional-Programs.pdf](https://caate.net/wp-content/uploads/2018/02/2020-Standards-Crosswalk_Final_for-Professional-Programs.pdf). Accessed July 31, 2018.
21. Practice analysis, 7th edition. Board of Certification, Inc Web site. [http://www.bocatac.org/system/document\\_versions/versions/24/original/boc-pa7-content-outline-20170612.pdf?1497279231](http://www.bocatac.org/system/document_versions/versions/24/original/boc-pa7-content-outline-20170612.pdf?1497279231). Accessed July 31, 2018.
22. National Athletic Trainers' Association. *Athletic Training Education Competencies*. 5th ed. Dallas, TX: National Athletic Trainers' Association; 2011.
23. Cowie MR, Blomster JJ, Curtis LH, et al. Electronic health records to facilitate clinical research. *Clin Res Cardiol*. 2017;106(1):1–9.
24. Nottingham SL, Lam KC, Kasamatsu TM, Eppelheimer BL, Welch Bacon CE. Athletic trainers' reasons for and mechanics of documenting patient care: a report from the Athletic Training Practice-Based Research Network. *J Athl Train*. 2017;52(7):656–666.
25. Welch Bacon CE, Eppelheimer BL, Kasamatsu TM, Lam KC, Nottingham SL. Athletic trainers' perceptions of and barriers to patient care documentation: a report from the Athletic Training Practice-Based Research Network. *J Athl Train*. 2017;52(7):667–675.
26. Valovich McLeod TC, Kostishak N Jr, Anderson BE, Welch Bacon CE, Lam KC. Patient, injury, assessment, and treatment characteristics and return to play timelines after sport-related concussion: an investigation from the Athletic Training Practice-Based Research Network. *Clin J Sport Med*. 2019;29(4):298–305.
27. Kerr ZY, Dompier TP, Dalton SL, Miller SJ, Hayden R, Marshall SW. Methods and descriptive epidemiology of services provided by athletic trainers in high schools: The National Athletic Treatment, Injury and Outcomes Network Study. *J Athl Train*. 2015;50(12):1310–1318.
28. Verhaagen E, van Mechelen W. *Sports Injury Research*. New York, NY: Oxford University Press, Inc; 2010.
29. Rothman KJ, Greenland S, Lash TL. *Modern Epidemiology*. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2008.
30. Wilkerson GB, Denegar CR. Cohort study design: an underutilized approach for advancement of evidence-based and patient-centered practice in athletic training. *J Athl Train*. 2014;49(4):561–567.
31. Teel EF, Marshall SW, Shankar V, McCrea M, Guskiewicz KM. Predicting recovery patterns after sport-related concussion. *J Athl Train*. 2017;52(3):288–298.
32. Wasserman EB, Herzog MM, Collins CL, Morris SN, Marshall SW. Fundamentals of sports analytics. *Clin Sports Med*. 2018;37(3):387–400.
33. Engelen-van Melick N, van Cingel REH, van Tienen TG, Nijhuis-van der Sanden MWG. Functional performance 2–9 years after ACL reconstruction: cross-sectional comparison between athletes with bone-patellar tendon-bone, semitendinosus/gracilis and healthy controls. *Knee Surg Sports Traumatol Arthrosc*. 2017;25(5):1412–1423.
34. Gribble PA, Terada M, Beard MQ, et al. Prediction of lateral ankle sprains in football players based on clinical tests and body mass index. *Am J Sports Med*. 2016;44(2):460–467.
35. McCrory P, Meeuwisse WH, Dvorak J, et al. Consensus statement on concussion in sport - the 5th International Conference on Concussion in Sport held in Berlin, October 2016. *Br J Sports Med*. 2018;51(11):838–847.
36. Portney LG, Watkins MP. *Foundations of Clinical Research*. 3rd ed. Upper Saddle River, NJ: Pearson Education, Inc; 2009.
37. Maffet M, Roton J Jr. Hemophilia in sports: a case report and prophylactic protocol. *J Athl Train*. 2017;52(1):65–70.
38. Swedler DI, Nuwer JM, Nazarov A, Huo SC, Malevanchik L. Incidence and descriptive epidemiology of injuries to college ultimate players. *J Athl Train*. 2015;50(4):419–425.
39. Lam KC, Snyder Valier AR, Valovich McLeod TC. Injury and treatment characteristics of sport-specific injuries sustained in interscholastic athletics: a report from the Athletic Training Practice-Based Research Network. *Sports Health*. 2015;7(1):67–74.
40. Lam KC, Valier AR, Anderson BE, McLeod TC. Athletic training services during daily patient encounters: a report from the Athletic Training Practice-Based Research Network. *J Athl Train*. 2016;51(6):435–441.
41. Marshall AN, Kikugawa TM, Lam KC. Patient, treatment, and cost characteristics associated with sport-related ankle sprains: a report from the Athletic Training Practice-Based Research Network [published online ahead of print July 30, 2019]. *Athl Train Sports Health Care*. doi: 10.3928/19425864-20190521-01
42. Simon JE, Wikstrom EA, Grooms DR, Docherty CL, Dompier TP, Kerr ZY. Athletic training service characteristics for patients with ankle sprains sustained during high school athletics. *J Athl Train*. 2019;54(6):676–683.
43. Grooms DR, Simon JE, Dalton SL, Dompier TP, Kerr ZY. High school athletic trainer services for knee injuries. *J Athl Train*. 2018;53(10):956–964.
44. Almquist J, Valovich McLeod TC, Cavanna A, et al. Summary statement: appropriate medical care for the secondary school-aged athlete. *J Athl Train*. 2008;43(4):416–427.
45. Lopes Sauers AD, Sauers EL, Snyder Valier AR. Quality improvement in athletic health care. *J Athl Train*. 2017;52(11):1070–1078.
46. Shanley E, Thigpen CA, Chapman CG, Thorpe J, Gilliland RG, Sease WF. Athletic trainers' effect on population health: improving access to and quality of care. *J Athl Train*. 2019;54(2):124–132.

47. Bachmann LM, Kolb E, Koller MT, Steurer J, ter Riet G. Accuracy of Ottawa ankle rules to exclude fractures of the ankle and mid-foot: systematic review. *BMJ*. 2003;326(7386):417.
48. Langley GJ, Moen RD, Nolan KM, Nolan TW, Norman CL, Provost LP. *The Improvement Guide: A Practical Approach to Enhancing Organizational Performance*. 2nd ed. San Francisco, CA: Jossey-Bass; 2009.
49. Vimalachandran P, Wang H, Zhang Y, Heyward B, Whittaker F. Ensuring data integrity in electronic health records: a quality health care implication. Paper presented at: International Conference on Orange Technologies; October 23–26, 2018; Melbourne, Australia.
50. Harrington L, Porch L, Acosta K, Wilkens K. Realizing electronic medical record benefits: an easy-to-do usability study. *J Nurs Adm*. 2011;41(7–8):331–335.
51. Penoyer DA, Cortelyou-Ward KH, Noblin AM, et al. Use of electronic health record documentation by healthcare workers in an acute care hospital system. *J Healthc Manag*. 2014;59(2):130–144.
52. Yee T, Needleman K, Pearson M, Parkerton P, Parkerton M, Wolstein J. The influence of integrated electronic medical records and computerized nursing notes on nurses' time spent in documentation. *Comput Inform Nurs*. 2012;30(6):287–292.
53. Christino MA, Matson AP, Fischer SA, Reinert SE, DiGiovanni CW, Fadale PD. Paperwork versus patient care: a nationwide survey of residents' perceptions of clinical documentation requirements and patient care. *J Grad Med Educ*. 2013;5(4):600–604.
54. Whittaker AA, Aufdenkamp M, Tinley S. Barriers and facilitators to electronic documentation in a rural hospital. *J Nurs Scholarsh*. 2009;41(3):293–300.
55. Zierler-Brown S, Brown TR, Chen D, Blackburn RW. Clinical documentation for patient care: models, concepts, and liability considerations for pharmacists. *Am J Health Syst Pharm*. 2007;64(17):1851–1858.
56. Welch Bacon CE, Kasamatsu TM, Lam KC, Nottingham SL. Future strategies to enhance patient care documentation among athletic trainers: a report from the Athletic Training Practice-Based Research Network. *J Athl Train*. 2018;53(6):619–626.
57. Lo W. Document like this, not that: CDI insights from the physician and CDI specialist perspective. *J AHIMA*. 2014;85(7):36–40.
58. Boling MC, Padua DA, Marshall SW, Guskiewicz K, Pyne S, Beutler A. A prospective investigation of biomechanical risk factors for patellofemoral pain syndrome: the Joint Undertaking to Monitor and Prevent ACL Injury (JUMP-ACL) cohort. *Am J Sports Med*. 2009;37(11):2108–2116.
59. Broglio SP, McCrea M, McAllister T, et al. A national study on the effects of concussion in collegiate athletes and US military service academy members: the NCAA-DoD Concussion Assessment, Research and Education (CARE) Consortium structure and methods. *Sports Med*. 2017;47(7):1437–1451.
60. Job settings: where ATs work. National Athletic Trainers' Association Web site. <https://www.nata.org/about/athletic-training/job-settings>. Published 2019. Accessed February 26, 2019.

---

*Address correspondence to Ashley N. Marshall, PhD, ATC, Department of Health and Exercise Science, Appalachian State University, ASU Box 32071, 1179 State Farm Road, Boone, NC 28608. Address e-mail to [marshallan1@appstate.edu](mailto:marshallan1@appstate.edu).*